



# Response of Different Panchagavya Levels on Yield Attributes of Bitter Gourd (*Momordica charantia* L.)

Rahul Yadav <sup>a++</sup>, Urfi Fatmi <sup>a#</sup>, Bhagchand Yadav <sup>b†\*</sup>,  
Udai Pal Singh <sup>b‡</sup> and Deepanshu <sup>a#</sup>

<sup>a</sup> Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, India.

<sup>b</sup> Directorate of Research, SKRAU, Bikaner, Rajasthan, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i84891>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120759>

Original Research Article

Received: 22/05/2024

Accepted: 24/07/2024

Published: 07/08/2024

## ABSTRACT

The field experiment on response of different panchgavya levels on yield attributes of bitter gourd (*Momordica charantia* L.) was conducted at Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj from January to July, 2023. The experiment was laid out in Factorial Randomized Block Design (FRBD) with three replications and eighteen treatment combination. The experiment consist of two factors. The factor one is hybrids as (TMBI-1309, TMBI-3346, VNR22) and factor second is different concentration of Panchagavya in which control (water spray), 1%

<sup>++</sup> Research Scholar;

<sup>#</sup> Assistant Professor;

<sup>†</sup> Senior Research Fellow;

<sup>‡</sup> Young Professional-II;

\*Corresponding author: E-mail: [bcyadavhorti@gmail.com](mailto:bcyadavhorti@gmail.com);

**Cite as:** Yadav, Rahul, Urfi Fatmi, Bhagchand Yadav, Udai Pal Singh, and Deepanshu. 2024. "Response of Different Panchagavya Levels on Yield Attributes of Bitter Gourd (*Momordica Charantia* L.)". *International Journal of Plant & Soil Science* 36 (8):618-23. <https://doi.org/10.9734/ijpss/2024/v36i84891>.

spray of Panchagavya, 2% spray of Panchagavya, 3% spray of Panchagavya, 4% spray of Panchagavya and 5% spray of Panchagavya. TMBI-3346 was recorded significantly higher yield attributes of bitter gourd over other hybrids in parameters like fruit weight (99.91 g), fruit length (15.64 cm), number of fruits per plant (37.0), fruit yield per plant (3.73 kg) and fruit yield (331.42 q/ha). Among The different Panchagavya treatments, 5% spray of Panchagavya resulted in significantly better fruit weight (104.66 g), fruit length (15.92 cm), number of fruits per plant (37.2), fruit yield per plant (3.90 kg) and fruit yield (346.99 q/ha). The interaction effect of TMBI-3346 + 5% spray of panchagavya resulted in significantly higher yield parameters.

**Keywords:** Bitter gourd hybrids; Panchagavya; bitter gourd.

## 1. INTRODUCTION

“Bitter gourd (*Momordica charantia* L.), is an important vegetable crop and is grown for its immature tuberculate fruits which have a unique bitter taste. It belongs to the family Cucurbitaceae. It is also known as bitter melon, bitter cucumber, balsam-pear, bitter apple, or bitter squash. This vegetable is also called karela in India, nigauri in Japan, goya in Okinawa, ampalaya in the Philippines, and ku-gua throughout China. *Momordica charantia* likely originated in eastern India or southern China. Today, it is found growing in fields across Asia, though it has also become popular in the Caribbean and South America. There are several types of bitter gourd, but the two most common are Chinese bitter gourd and Indian bitter gourd. Fruits are considered to be a rich source of vitamins and minerals and it contains 88 mg vitamin C per 100 g. Bitter gourd is rich in a number of important antioxidants as well. In fact, half a cup of fresh bitter gourd accounts for about 93% of recommended daily intake of vitamin C. The younger the fruit, the more vitamin C it contains. Bitter gourd has been used as a folk remedy for tumors, asthma, skin infections, gastro intestinal problems” [1-3].

Bitter gourd survives in warm and humid climates, typically in temperatures required from 24°C to 30°C (75°F to 86°F). The ideal soil for its cultivation is well-drained loamy soil with a pH level around 6.0 to 7.0. As bitter gourd plants spread quite vigorously nutrient management becomes crucial for optimal yield [4]. The use of organic and inorganic nutrients increased fruit length, diameter, weight, number and yield of bitter gourd and nowadays more emphasis is given to nutrients of organic region, among which Panchagavya is a potential nutrient supplier.

“Panchagavya is a term used in Ayurveda for fermented product made from five ingredients


obtained from cow such as milk, urine, dung, curd and butter” [1]. “In Sanskrit, all these five products are individually called “Gavya” and collectively termed as Panchagavya. The panchagavya is an efficient plant growth stimulant that enhances the biological efficiency of crops. It is used to activate soil and to protect the plants from diseases and also increase the nutritional quality of fruits and vegetables. It may be used as a foliar spray, as soil application along with irrigation water, seed or seedling treatment etc. Three per cent panchagavya is an ideal concentration for the foliar spray” [5]. “Bio-chemical properties of panchagavya revealed that it possesses almost all the major nutrients like N, P, K and micro nutrients essential for plant and growth hormones like IAA and GA required for crop growth. Role of foliar applied panchagavya in production of many plantations crops has been well documented in India” [6]. “Panchagavya is a popular foliar nutrition prepared by organic growers of Tamil Nadu as an indigenous material and used widely for agricultural and horticultural crops” [7].

## 2. MATERIALS METHODS

The field experiment was conducted in the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during January to July, 2023. The experiment was laid out in Factorial Randomized Block Design (FRBD) with three replications. The experiment consisted of two factors Factor A: (TMBI-1309, TMBI-3346, VNR-22) and six levels of Panchagavya spray *i.e.*, (Factor B) 0 %, 1 %, 2 %, 3%, 4 %, 5%. In each hill had one plant at a spacing of 150 cm x 75 cm.

## 3. RESULTS AND DISCUSSION

The data on yield attributes of bitter gourd after application of Panchagavya was recorded and is presented in Table 1 and illustrated through



**Fig. 1. Effect of panchagavya on yield (q/ha) of different hybrids of bitter gourd**

**Table 1. Effect of panchagavya on fruit of different hybrids of bitter gourd**

Treatment	Fruit Weight (g)				Fruit Length (cm)				Number of fruits Per Plant			
	TMBI-1309	TMBI-3346	VNR-22 (check)	Mean	TMBI-1309	TMBI-3346	VNR-22 (check)	Mean	TMBI-1309	TMBI-3346	VNR-22 (check)	Mean
Water control	80.67	87.49	82.48	83.55	12.55	13.80	12.92	13.09	27.4	32.5	30.6	30.2
1% PG	82.65	91.54	88.74	87.64	12.96	14.69	14.21	13.95	28.9	34.0	32.9	32.0
2% PG	88.74	98.18	92.82	93.25	13.71	15.64	14.54	14.63	30.8	36.3	34.0	33.7
3% PG	92.31	103.62	98.75	98.23	13.94	16.01	15.26	15.07	32.7	39.1	36.5	36.1
4% PG	97.98	108.37	102.49	102.95	14.60	16.71	15.53	15.61	33.9	39.8	37.0	36.9
5% PG	99.99	110.27	103.71	104.66	15.01	16.96	15.78	15.92	34.0	40.5	37.1	37.2
<b>Mean</b>	90.39	99.91	94.83		13.79	15.64	14.71		31.3	37.0	34.7	
<b>Factors</b>	<b>F Test</b>	<b>SE(d)</b>	<b>CD<sub>0.05</sub></b>		<b>F Test</b>	<b>SE(d)</b>	<b>CD<sub>0.05</sub></b>		<b>F Test</b>	<b>SE(d)</b>	<b>CDS<sub>0.05</sub></b>	
<b>Factor H</b>	S	1.739	3.669		S	0.344	0.726		S	0.839	1.770	
<b>Factor B PG</b>	S	2.459	5.188		S	0.487	1.027		S	1.186	2.503	
<b>H x PG</b>	S	4.259	8.986		S	0.843	1.778		S	2.055	4.336	

PG = Panchgavya; H= Hybrids

**Table 2. Effect of panchagavya on yield of different hybrids of bitter gourd**

Treatment	Yield Per Plant (kg)				Yield (q/ha)			
	TMBI-1309	TMBI-3346	VNR-22 (check)	Mean	TMBI-1309	TMBI-3346	VNR-22 (check)	Mean
Water control	2.21	2.84	2.52	2.52	196.69	252.55	224.00	224.41
1% PG	2.39	3.12	2.92	2.81	212.62	277.12	259.45	249.73
2% PG	2.76	3.59	3.18	3.18	244.87	318.97	282.91	282.25
3% PG	3.02	4.05	3.60	3.56	268.15	359.93	320.08	316.05
4% PG	3.33	4.31	3.79	3.81	295.62	383.28	336.90	338.60
5% PG	3.40	4.46	3.85	3.90	302.40	396.66	341.92	346.99
<b>Mean</b>	2.85	3.73	3.31		253.39	331.42	294.21	
<b>Factors</b>	<b>F Test</b>	<b>SE(d)</b>	<b>CD<sub>0.05</sub></b>		<b>F Test</b>	<b>SE(d)</b>	<b>CD<sub>0.05</sub></b>	
<b>Factor H</b>	S	0.134	0.283		S	11.902	25.112	
<b>Factor B PG</b>	S	0.189	0.400		S	16.832	35.513	
<b>H x PG</b>	S	0.328	0.692		S	29.155	61.511	

Fig 1. Among the treatments with respect to the yield attributes of bitter gourd. Yield attributes of bitter gourd varies significantly between different hybrids and Panchgavya treatments. Hybrid TMBI-3346 recorded significantly higher fruit weight (99.91 g), fruit length (15.6cm), number of fruits per plant (37.0), fruits yield per plant (3.73 kg) and fruit yield per hectare (331.42 q/ha) and the lower in TMBI-1309 hybrid of fruit weight (90.4g), fruit length (13.79cm), number of fruits per plant (31.3), fruits yield per plant (2.85 kg) and fruit yield per hectare (253.4 q/ha) were recorded in hybrid TMBI-1309. Among the panchgavya treatments 5% spray result in the higher fruit weight (104.66 g), fruit length (15.92 cm), number of fruits per plant (37.2), fruit yield per plant (3.90 kg), and fruit yield per hectare (346.99 q/ha) while lower fruit weight (83.5g), fruit length (13.09cm), number of fruits per plant (30.2), fruits yield per plant(2.52kg) and fruit yield per hectare (224.41q/ha) were recorded in 5 % spray of Panchagavya.

The interaction effect of hybrids and Panchgavya levels resulted in significant difference among the treatments. higher fruit weight of (110.27 g) was reported in hybrids TMBI-3346 with 5% Panchgavya spray which was found to be at par with treatment combination VNR-22 + 4% Panchgavya spray (102.5 g) and VNR-22 + 5% Panchgavya spray (103.7 g), while as lower weight of (80.67 g) was reported in TMBI-1309 with 0% Panchgavya spray. and higher fruit length of (16.96 cm) was reported in hybrids TMBI-3346 with 5% Panchgavya spray which was found to be at par with treatment combination TMBI 3346 + 4% Panchgavya spray (16.71 cm) and TMBI-3346 + 3% Panchgavya spray (16.01 cm) and VNR-22 + 5% Panchgavya spray (15.78 cm) and VNR-22 + 4% Panchgavya spray (15.53.cm) while as lower fruit length of (12.55 cm) was reported in TMBI-1309 with 0% Panchgavya spray. and higher number of fruits per plant (40.5) was reported in hybrids TMBI-3346 with 5% Panchgavya spray which was found to be at par with treatment combination TMBI 3346 + 4% Panchgavya spray (39.8) and TMBI-3346 + 3% Panchgavya spray (39.1) and VNR-22 + 5% Panchgavya spray (37.1) and VNR-22 + 4% Panchgavya spray (37.5) while as lesser number of fruits per plant (27.4) was reported in TMBI-1309 with 0% Panchgavya spray. and higher yield per plant (4.46 kg) was reported in hybrids TMBI-3346 with 5% Panchgavya spray which was found to be at par with treatment

combination TMBI 3346 + 4% Panchgavya spray (4.31 kg) and TMBI-3346 + 3% Panchgavya spray (4.05 kg) and VNR-22 + 5% Panchgavya spray (3.85 kg) and VNR-22 + 4% Panchgavya spray (3.79) while as lesser yiled per plant (2.21 kg) was reported in TMBI-1309 with 0% Panchgavya spray. and higher yield quintal per hactare (396.66 q/ha) was reported in hybrids TMBI-3346 with 5% Panchgavya spray which was found to be at par with treatment combination TMBI 3346 + 4% Panchgavya spray (383.28 q/ha) and TMBI-3346 + 3% Panchgavya spray (359.93 q/ha) and VNR-22 + 5% Panchgavya spray (341.92 q/ha) and VNR-22 + 4% Panchgavya spray (336.9 q/ha) while as lesser yiled per plant (196.69) was reported in TMBI-1309 with 0% Panchgavya spray.

Significantly more number of fruits per plant was recorded with 5% spray of Panchagavya which might be due to presence of auxin and kinetin in Panchagavya which upon application might have favoured the plants to produce more number of fruits per plant Significant and longer vine was recorded with the application of 5% spray Panchagavya which may be due to its optimum concentration as it contains auxin which leads to apical dominance and gibberellic acid that leads to cell elongation and cell division resulting in increased vin length These findings are in close conformity with those of Gore and Sreenivasa [3] in tomato (*Lycopersicon esculentum* L. Mill.) Sivakumar [8], Swain et al. [9] in chilli (*Capsicum annum* L.) cv. Kuchinda local. Significant and higher fruit yield was recorded with application of 5% spray of Panchagavya which might be due to microbes present in Panchagavya that produces growth hormones which may cause increase in fruit weight, number of fruits/plant. These findings are in close conformity with those of Boraiah et al. [2] and Suchitra et al. [10]. In capsicum (*Capsicum annum* L. var *grossum*) and *Abelmoschus esculentus* L.

#### 4. CONCLUSION

Based on the results of the present investigation entitled "Response of Different Panchagavya Levels on Yield Attributes of Bitter Gourd (*Momordica charantia* L.)" it is concluded that among the different treatment combination of hybrids and Panchagavya levels TMBI-3346 + 5% spray of Panchagavya Performed significantly better in yield attributes of bitter gourd.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Amalraj VA, Rakkiyappan, Devi, Reema AK. Evaluation of wild sugarcane erianthus Arudinaceus (Retz) Jesw. Germplasm. Journal of Sugarcane Research. 2011;1(2):23-27.
2. Boraiah B, Devkumar N, Shubha S, Palanna KB. Effect of panchagavya, jeevnamrutha and cow urine on beneficial microorganisms and yield of capsicum (*Capsicum annum L. var grossum*). Int. J. Curr. Microbiol. App. Sci. 2017;6(9):3226-3234.
3. Gore N, Sreenivasa MN. Influence of liquid organic manures on growth, nutrient content and yield of tomato (*Lycopersicon esculentum L. Mill.*) in the sterilized soil. Karnataka Journal of Agricultural Science. 2011;2:153-157.
4. Mohan Kumar K. Influence of Various Organic Inputs on Growth and Yield of Snake Gourd (*Trichosanthes anguina L.*). International Journal of Agriculture Sciences; 2016. ISSN. 2016:0975-3710.
5. Sarkar S, Kundu SS, Ghoria D. Validation of ancient liquid organics- Panchagavya kunapajala as plant growth promoters. Indian Journal of Traditional Knowledge. 2014;13(2):398-403.
6. Selvaraj N. Report on the work done on organic farming at Horticultural Research Station, Ooty, Tamil Nadu Agric. Univ. 2003;2-5.
7. Swaminathan C, Swaminathan V, Vijayalakshmi V. Panchagavya boon to organic farming. International book Distributing Corporation, India; 2007.
8. Sivakumar T. Review on panchagavya. Int. J. Adv. Res. Biol. Sci. 2014;1(8):130-154.
9. Swain SS, Sahu GS, Mishra N. Effect of panchagavya on growth and yield of chilli (*Capsicum annum L.*) cv. Kuchinda local. Green Farming. 2015;6(2):338-340.
10. Suchitra S, Poonguzhali S, Suguna S, Jothibas K. Effect of panchagavya on growth and yield of Abelmoschus esculentus L. cv. Arka Anamika. Int. J. Curr. Microbiol. App. Sci. 2017;6(9):3090-3097.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/120759>