



***In vivo* Hepatoprotective and Antioxidant Activity of *Enicostemma littorale* against CCl₄ Induced Liver Damage in Rats**

Dhanapal Venkatachalam^{1*}, B. Samuel Thavamani¹, K. Muddukrishniah¹ and Sukanya Vijayan¹

¹*Department of Pharmacognosy, Sanjo College of Pharmaceutical Studies, Velappara, Palakkad, Kerala – 678 702, India.*

Authors' contributions

This work was carried out in collaboration between all authors. Author DV designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors BST and KM managed the analyses of the study. Author SV managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRIMPS/2018/43512

Editor(s):

(1) Dr. John Yahya I. Elshimali, Professor, Department of Pathology and Oncology, UCLA School of Medicine, Charles R. Drew University of Medicine and Science, California, USA.

Reviewers:

- (1) Muhammad Shahzad Aslam, Pakistan.
(2) G. Bupesh, Bharath Institute of Higher Education & Research, Sree Balaji Medical College & Hospital, India.
(3) T. Nagamma, Department of Biochemistry, Melaka Manipal Medical College, India.

Complete Peer review History: <http://www.sciencedomain.org/review-history/26236>

Original Research Article

Received 15 June 2018
Accepted 01 September 2018
Published 14 September 2018

ABSTRACT

To investigate the hepatoprotective and antioxidant effects of ethanol extract of *Enicostemma littorale* Blume (Ens) against CCl₄ induced hepatic injury in albino rats. Male albino rats of six numbers in each group were undertaken for study. Hepatoprotective and antioxidant effect of *E. littorale* Blume (Ens) ethanol extract at a dosage of 100 & 200 mg/kg body weight was evaluated. The degree of hepatoprotection was assessed by measuring the activity levels of the marker enzymes such as serum aspartate transaminase (AST), alanine transaminase (ALT) alkaline phosphatase (ALP), acid phosphatase (ACP) and total bilirubin. Lipid peroxidation caused by free radicals was assessed by measuring the activity levels of the tissue antioxidant enzymes such as glutathione peroxidase (GPX), catalase (CAT), superoxide dismutase (SOD). The CCl₄

*Corresponding author: E-mail: vddpaul@gmail.com;

administered rats recorded elevated activity levels of serum AST, ALT, ALP and ACP revealing CCl₄ induced hepatotoxicity. In the groups treated with 100 mg/kg and 200 mg/kg of the extract, the above biochemical markers of hepatotoxicity were found to be decreased when compared to CCl₄ treated control group. Both the doses of ethanol extract of *Enicostemma littorale* (EEEL) used in the study showed significant protective property than the control. In the groups treated with 100 mg/kg and 200 mg/kg of the extract, GPX, SOD and catalase were found to be increased when compared to CCl₄ treated control group. It can be concluded that the ethanol extract of *E. littorale* Blume is not only hepatoprotective but also possess significant antioxidant property.

Keywords: Hepatoprotective; antioxidant enzymes; *Enicostemma littorale* Blume.

1. INTRODUCTION

The hepatic system is a very vital organ system involved in the body's metabolic activities. Hence, the chemical reactions in the liver may generate several reactive species like free radicals. These reactive species form covalent bond with the lipids of the tissue. However inbuilt protective mechanisms combat the hazardous reactions associated with the free radicals. Due to excessive exposure to hazardous chemicals, the free radicals generated will be so high such that they overpower the natural defensive system leading to hepatic damage and cause jaundice, cirrhosis and fatty liver, which remains one of the serious health problems. Carbon tetrachloride (CCl₄) is one such hazardous chemical which induces hepatopathy through membrane lipid peroxidation by its free radical derivative like CCl₃·, CCl₃O₂·. Excessive production of the reactive species manifests in tissue-thiol depletion, lipid peroxidation, plasma membrane damage etc., culminating into severe hepatic injury [1]. Much of the cell damage that occurs during liver degeneration is believed to be caused by free radicals which are highly reactive oxygen species liberated during alcohol metabolism. These radicals react with cell membrane and induce lipid peroxidation, which has been implicated as an important pathological mediation [2] in many diseases such as heart disease, diabetes, cancer and liver disease. Management of liver diseases is still a challenge to modern medicine. In such a background, it is realized that antioxidant activity or inhibition of generation of free radicals plays a crucial role in providing protection against such hepatic damage. Several herbs and herbal products are known to possess antioxidant constituents and may be used as organ protective agents.

Plant drugs are frequently considered to be less toxic and free from side effects [3]. Numerous medicinal plants and their formulations are used for liver disorders in ethnomedical practices as

well as in traditional systems of medicine in India. There are a number of evidences indicating that natural substances from edible and medicinal plants exhibit strong antioxidative activity, and could work against hepatic toxicity caused by various toxicants [4-5] *Enicostemma littorale* (*E. littorale*) Blume (Family: Gentianaceae) is a perennial, tropical traditional medicinal herb with sessile lanceolate leaves, flowers arranged in clusters and fruit in a capsule. The plant is locally used for its medicinal properties such as anti-inflammatory, antiulcer activity [6], hypoglycaemic [7] and antimalarial activities [8] in Tamilnadu and Kerala, India. The present study aims to investigate the hepatoprotective and antioxidant effects of *E. littorale* on CCl₄ induced hepatotoxicity in male albino rats.

2. MATERIALS AND METHODS

2.1 Plant Material

The whole plant of *E. littorale* was collected during January (2018) in and around Palakkad District, Kerala, India. The plant was cleaned, shade dried and authenticated by Dr. Jayaraman, Plant Anatomy Research Centre, Chennai. A voucher specimen was deposited in the Department of Pharmacognosy, Sanjo College of Pharmaceutical Studies, Palakkad, Kerala (SCOPS/P.COG/07/2018).

2.2 Preparation of Plant Extract

For preliminary phytochemical analysis, extract was prepared by weighing 500 grams of the dried powdered leaves and subjected to hot successive continuous extraction using Soxhlet apparatus with different solvents like petroleum ether, benzene, chloroform, ethanol and water in the increasing order of polarity. The extracts were filtered in each step using Whatman filter paper. The filtrate was concentrated using a rotary evaporator at low temperature (40-45°C)

and pressure. The presence or absence of the primary and secondary phytoconstituents was detected by usual prescribed method [9].

2.3 Chemicals and Drugs

Carbon tetra chloride and Ecoline kits for serum aspartate transaminase (AST), alanine transaminase (ALT), alkaline phosphatase (ALP), acid phosphatase (ACP) and total bilirubin, were purchased from Sigma Co. (Sigma St. Louis, MO). Standard Silymarin was obtained from Ranbaxy Ltd, New Delhi. Absolute ethanol was of analytical grade and was purchased from Merck (German). The other reagents were of analytical grade.

2.4 Animals

Wistar albino rats of 180-230 gm, maintained in the Animal House facility of the Department of Pharmacology, Sanjo College of Pharmaceutical Studies, Palakkad were used in these experiments. The animals were fed on standard small animal feeds (Excel feed, Ilorin) and water *ad libitum*. This research was carried out by following the rules governing the use of laboratory animals as accepted internationally. The experiment was conducted between the hours of 900 h and 1600 h. The experimental groups consisted of six animals. They were maintained at constant room temperature ($22^{\circ} \pm 1^{\circ}\text{C}$) and submitted to 12 h light/dark cycle with free access to food and water.

2.5 Experimental Procedure

2.5.1 Acute oral toxicity study

Acute oral toxicity was conducted as per OECD guidelines (Organization of Economic Cooperation and Development) 423 (Acute toxic class method). The acute toxic class method is a step wise procedure of three animals of single sex per step. Depending on the mortality and / or moribund status of animals, on an average 2-4 steps may be necessary to allow judgment on the acute toxicity of the test substance. This procedure results in the use of a minimal number of animals while allowing for acceptable data based scientific conclusion. The method uses defined doses (5, 50, 300, 2000 mg/kg body weight) and the results allow a substance to be ranked and classified according to the Globally Harmonized System (GHS) for the classification of chemicals which causes acute toxicity. The method previously described by Lorke [10] was adopted. The LD50 was found to be 2000 mg/kg body weight.

2.6 Hepatoprotective Activity

The method of Ko et al. [11] was used for screening the hepatoprotectivity of the ethanol extract of *Enicostemma littorale* prepared as mentioned above. Adult Wistar rats of either sex were randomly assigned into 5 groups of 6 animals each. The animals of Group I served as normal control and received only the vehicle normal saline (10 ml/kg i.p). Group II served as toxic control and administered CCl_4 (1 ml/kg) by subcutaneous injection. The animals of Group III and IV received *Enicostemma littorale* extract (100 mg/kg BW and 200 mg/kg BW p.o. respectively) for 15 days. Group V served as Standard and was treated with Silymarin (25 mg/kg BW i.p., for 15 days). The animals (except Group 1) were treated CCl_4 at a dose of 1 ml/kg BW by subcutaneous injection. Blood samples were collected after last dose of CCl_4 administration by direct cardiac puncture under light ether anesthesia and animals were sacrificed by cervical decapitation and hepatic tissue was collected. Heparinized blood sample was taken and assessed for serum enzyme markers and hepatic tissue was taken and subjected to histopathological study and further tissue was analyzed for Glutathione and lipid peroxidation.

- Group 1-Normal animals (10 mg/kg i.p)
- Group 2- CCl_4 (1 ml/kg s.c) treated animals
- Group 3- CCl_4 + EEEL (100 mg/kg, p.o.) treated animals.
- Group 4- CCl_4 + EEEL (200 mg/kg, p.o.) treated animals.
- Group 5- CCl_4 + Silymarin (25 mg/kg i.p.) treated animals

2.7 Estimation of Biochemical Parameters

Serum aspartate transaminase (AST), alanine transaminase (ALT) alkaline phosphatase (ALP) and acid phosphatase (ACP), and total bilirubin were estimated in the separated serum [12]. The tissue levels of enzymatic antioxidants viz. superoxidase dismutase (SOD), catalase (CAT), glutathione peroxidase (GPX), were also estimated [13-16].

2.8 Estimation of Glutathione Peroxidase, Superoxide Dismutase (SOD), and Catalase (CAT)

Glutathione peroxidase (GPx) was measured by the method described by Rotruck et al. The

activities were expressed as μg of GSH consumed/ min/mg protein. The SOD activity in cell lysate was determined as per the method followed earlier by Kakkar et al. Enzyme activity was expressed as 1 unit = 50% inhibition/minute/mg of protein. The CAT activity in cell lysate was assayed as per the method reported earlier by Aebi. CAT was expressed in terms of μmol of hydrogen peroxide decomposed/min/mg of protein [17-19].

2.9 Statistical Analysis

The results are analysed by one way ANOVA followed by Dunnett's test using prism pad software and p value <0.001 was considered to be significant.

3. RESULTS

3.1 Acute Toxicity

The acute toxicity in Wistar albino rat was studied as per the Organization for Economic Cooperation and Development (OECD) guideline 423. The LD₅₀ value of EEEL i.e 2000 mg/kg body weight was determined using the method of maximum likelihood.

3.2 Hepatoprotective and Antioxidant Activity

The estimated values of serum AST, ALT, ALP, ACP and Total bilirubin values in control (saline) group of rats are tabulated in Table 1. A remarkable elevation was observed in Serum AST, ALT, ALP, ACP, total bilirubin and values in CCl₄ intoxicated rats (Toxic Control group). In the groups treated with 100 mg/kg and 200 mg/kg of the extract, the above bio-chemical markers of hepatotoxicity were found to be decreased when compared to CCl₄ treated control group. Evidently, the hepatoprotective effects of higher

dose of Ethanolic extracts of *Enicostemma littorale* (200 mg/kg) were near to that of standard i.e. Silymarin (25 mg/kg). Both the doses of EEEL used in the study showed significant protective property than control (*p<0.01, **p<0.001 vs. control). However, the test extract was found to be less potent than that of standard.

The tissue glutathione was found to be depleted upon CCl₄ intoxication which indicates that the tissue damage is due to over powering the inbuilt free radical scavenger mechanisms. This tissue Glutathione (GSH) depletion was inhibited by the pretreatment with test extract in a dose dependant manner. Similarly, lipid peroxidation induced by CCl₄ treatment was reversed by test extract in a dose dependant manner. The results are compiled in Table 2. In the groups treated with 100 mg/kg and 200 mg/kg of the extract, GPX, SOD and catalase were found to be increased when compared to CCl₄ treated control group. Evidently, the hepatoprotective effects of higher dose of Ethanolic extracts of *Enicostemma littorale* (200 mg/kg) were near to that of standard i.e. Silymarin (25 mg/kg). Both the doses of EEEL used in the study showed significant protective property than control. (*p<0.01, vs. control) However the test extract was found to be less potent than that of standard drug.

4. DISCUSSION

Since the extract has demonstrated for antioxidant property in dose dependant in all the models of the study, the ethanol extract was taken for assessing the *in vivo* hepatoprotective properties. Pretreatment with the test extract has reduced the elevated levels of biochemical markers of hepatotoxicity. Further it was also observed that the depletion of GSH in the issue was due to CCl₄ challenge and was reversed by

Table 1. Effect of EEEL on serum biochemical parameters

Treatment	AST(U/I)	ALT(U/I)	ALP (IU/L)	ACP(U/L)	Total Bilirubin (mg/100 ml of blood)
Control (saline)	97.3±1.18	35.08±0.2	15.92±0.72	10.5±0.064	0.39±0.04
CCl ₄ (1 ml/kg)	186.7±1.82	136.9±1.94	98.3±7.9	38.6±2.9	0.89±0.76
EEEL(100 mg/kg)	138.64±5.92	72.4±6.8	63.9±5.8	25.4±0.95	0.74±0.06
EEEL(200 mg/kg)	106.8±8.28	47.5±4.1	43.6±3.4	16.5±0.18	0.57±0.04
Silymarin 25 mg/kg	105.3±4.3	49.4±3.6	34.8±2.9	16.2±1.2	0.24±0.03

Data are expressed as mean \pm S.E (n=6) *p<0.01 vs. control, **p<0.001 vs. control

EEEL- Ethanolic extract of *Enicostemma littorale*; AST-Aspartate transaminase; ACP-Acid phosphatase;

ALT- Alanine transaminase; ALP- Alkaline phosphatase




Fig. 1. Effect of EEEL on serum biochemical parameters

Table 2. Antioxidant activity of EEEL

Treatment	Glutathione peroxidase (mg liver protein) ⁻¹	SOD (mg liver protein) ⁻¹	Catalase (mg liver protein) ⁻¹
Control (saline)	0.992±0.05	75.81±1.94	296.83±10.05
CCl ₄ (1 ml/kg)	0.61±0.03	47.84±0.50	179.73±5.78
EEEL(100 mg/kg)	0.85*±0.07	67.73*±0.54	266.27*±8.74
EEEL(200 mg/kg)	0.92*±0.06	86.97*±0.75	281.38*±9.92
Silymarin 25 mg/kg	0.95*±0.03	88.34*±2.54	268.27*±6.46

Data are expressed as Mean ±S.E (n=6) *p<0.01 Vs control

EEEL- Ethanolic extract of *Enicostemma littorale*

SOD- Superoxidase dismutase




Fig. 2. Antioxidant activity of EEEL

the test extract and also reduced the extent of lipid peroxidation. Most of the mammals have an effective mechanism to prevent and neutralize the free radical induced damage, which is accomplished by a set of endogenous substances such as superoxide dismutase, catalase, and glutathione peroxidase and glutathione reductase. In biochemical system, superoxide radical and H_2O_2 react together to form the hydroxyl radical. This can attack and destroy almost all known biochemicals [20]. The hydroxyl radicals thus produced may attack the sugar of DNA base causing sugar fragmentation, base loss and DNA strand leakage. Ethanol extract of *E. littorale* reduced the superoxide anions and also scavenges off the hydroxyl radicals and hence, inhibit the cellular damage. It is apparent from the present study that the test extract does not interfere with the generation of the free radicals but it scavenges off the free radicals. CCl_4 undergoes hepatic metabolism to give rise to tri-chloromethyl radicals which, upon reacting with reactive oxygen species yields trichloromethyl peroxide radicals, which forms covalent bond with membrane lipids and destroy the membrane integrity. The observation of increased malondialdehyde (MDA) formation in hepatic cells after CCl_4 challenge is in accordance with the earlier report which suggests involvement of trichloromethyl and trichloromethylperoxy radicals in the propagation of per-oxidation process [21]. Pretreatment with the extract has prevented oxygen free radicals and thereby prevented the formation of peroxy radicals. This aspect of test extract also contributes to the hepatoprotectivity. Unpublished data on the hepatoprotective activity of this plant on other models like paracetamol and thiocetamide induced hepatotoxicity indicated that the hepatoprotectivity of the test extract is not model specific. SOD is metalloproteins catalyzing the dismutation of superoxide anion to hydrogen and oxygen. Studies have shown the importance of SOD in protecting cells against oxidative stress [22]. The SOD activity could be decreased in tissue during alcohol ingestion. This decrease could be due to the feedback inhibition or oxidative inactivation of enzyme protein due to excess ROS generation [23]. CAT, hemeprotein, catalyzes the reduction of hydrogen peroxides [24] acts as preventive antioxidant and plays an important role in protection of tissues against the deleterious effects of lipid peroxidation [25]. The activity levels of catalase in tissue decreased in ethanol fed animals. This might be due to the inhibition of CAT activity, which is suggestive of enhanced

synthesis of O_2^- during the ingestion of alcohol since O_2^- is a powerful inhibitor of catalase [26]. GPX is an enzyme with selenium in the form of selenocysteine and can catalyze the reduction of hydrogen peroxide and hydroperoxides to nontoxic products. GPX has a well-established role in protecting cells against oxidative injury. GPX is non-specific for H_2O_2 and lipid peroxide generated during alcohol ingestion which are efficiently be scavenged by GPX activity. The depression of this enzyme activity reflects perturbations in normal oxidative mechanism during alcohol ingestion. The cellular antioxidant defense enzymes viz. SOD, CAT, and GPX were significantly reduced in the CCl_4 administered rat. This might lead to decreased antioxidant defense and increased oxidative stress and thereby the tissue injury [27-29].

5. CONCLUSION

The results of the present investigation, concluded that the ethanol extract of the whole parts of *Encostemma littorale* possess significant hepatoprotective activity against carbon tetrachloride induced hepatotoxicity and antioxidant activity. The antioxidant potential may be attributed to the presence of polyphenolic compounds. Further studies like isolation and characterization of the active principle(s) responsible for such activity are needed to confirm.

CONSENT

It is not applicable.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Vir Ji Chrungoo, Kuldip Singh, Jaswant Singh. Differential biochemical response of freshly isolated rat hepatocytes to paracetamol, carbon tetrachloride and D-galactosamine toxicity. Ind J Exp Bio. 1997;35:603-610.

2. Muriel P. Role of free radicals in liver diseases. *Hepatol Int.* 2009;3:526–536.
3. Pari L, Saravanan R. Antidiabetic effect of diasulin, an herbal drug, on blood glucose, plasma insulin and hepatic enzymes of glucose metabolism in hyperglycaemic rats. *Diabetes Obes Metab.* 2004;6:286–292.
4. Upur H, Amat N, Blazekovic B, Talip A. Protective effect of *Cichorium glandulosum* root extract on carbon tetrachloride-induced and galactosamine induced hepatotoxicity in mice. *Food Chem Toxicol.* 2009;47:2022–2030.
5. Zeashan H, Amresh G, Singh S, Rao CV. Hepatoprotective activity of *Amaranthus spinosus* in experimental animals. *Food Chem Toxicol.* 2008;46:3417–3421.
6. Roy S, Niranjana C, Jyothi T, Shankrayya M, Vishwanath K, Prabhu K, et al. Antiulcer and anti-inflammatory activity of aerial parts *Encicostemma littorale* Blume. *J Young Pharm.* 2010;2(4):369–373.
7. Vishwakarma SL, Sonawane RD, Rajani M, Goyal RK. Evaluation of effect of aqueous extract of *Encicostemma littorale* Blume in streptozotocin induced type 1 diabetic rats. *Indian J Exp Biol.* 2010;48: 26–30.
8. Yghemonos S. The Earth Institute, University of Colombia. Malaria and poverty: European alliance against malaria. 2007;1–4.
9. Kokate CK. *Practical Pharmacognosy*, M/S Vallabh Prakashan, Pune. 1985;111-115.
10. Lorke D. A new approach to acute toxicity testing. *Archives of Toxicology.* 1983;54: 275-287.
11. Ko KM, Yick PK, Chiu TW, Hui TY, Cheng CHK, Kong YC. Impaired antioxidant status in CCl₄ intoxicated rats: An *in-vivo* study. *Fitotherapia* 1993;LXIV:539-544.
12. Malloy HT, Evelyn KA. The determination of bilirubin with the photoelectric colorimeter. *J of Biol Chem.* 1937;19:481-490.
13. Kakkar P, Das B, Viswanathan PN. A modified spectrophotometric assay of superoxide dismutase. *Indian J Biophys.* 1984;21:130–132.
14. Sinha AK. Colorimetric assay of catalase. *Anal Biochem.* 1972;47:389–394.
15. Rotruck JT, Pope AL, Ganther HE, Swason AB, Hafeman DG, Hoekstra WG. Selenium: Biochemical role as a component of glutathione peroxidase. *Science.* 1973;179:588–590.
16. Habig WH, Pabst MJ, Jakpoly WB. Glutathione-s-transferase: The first enzymatic step in mercapturic acid formation. *J Biol Chem.* 1974;249:7130–7139.
17. Rotruck JT, Pope AL, Ganther HE, Swanson AB, Hafeman DG, Hoekstra WG. Selenium: Biochemical role as a component of glutathione peroxidase. *Sci.* 1973;179:588-590.
18. Kakkar P, Das, Vishwanath PN. A modified spectrophotometric assay of superoxide dismutase. *Indian J Biochem Biophys.* 1984;21:130-132.
19. Aebi H. Catalase. In: Packer L, Ed. *Methods in Enzymatic Analysis*, Vol. II. New York: Academic Press. 1974;673-84.
20. Sasanka Chakraborty, Asha Naik S, Gali Reddy R. Phenylhydrazine mediated degradation of Bovine serum albumin and membrane proteins of human erythrocytes. *Biochem et Biophys Acta.* 1990;1028:89-94.
21. Indu Bala Koul, Aruna Kapil. Evaluation of the liver protective potential of Piperine, an active principle of black and long pep-pers. *Planta Med.* 1993;59:413-417.
22. Yang ES, Lee JH, Park JW. Ethanol induces peroxynitrite-mediated toxicity through inactivation of NADP⁺-dependent isocitrate dehydrogenase and superoxide dismutase. *Biochimie.* 2008;90(9):1316–1324.
23. Shanmugam KR, Ramakrishna CH, Mallikarjuna K, Reddy KS. Protective effect of ginger against alcohol induced renal damage and antioxidant enzymes in male albino rats. *Indian J Exp Biol.* 2010;48: 143–149.
24. Dash DK, Yeligar VC, Nayak SS, Ghosh T, Rajalingam D, Sengupta P, et al. Evaluation of hepatoprotective and antioxidant activity of *Ichnocarpus frutescens* (Linn.) R.Br. on paracetamol-induced hepatotoxicity in rats. *Trop J Pharm Res.* 2007;6(3):755–765.
25. Dinkova-Kostova AT. Protection against cancer by plant phenyl propenoids: induction of mammalian anticarcinogenic enzymes. *Mini Rev Med Chem.* 2002;2:595–610.
26. Popovic M, Janicijevic-Hudomal S, Kaurinovic B, Rasic J, Trivic S. Effects of various drugs on alcohol-induced oxidative stress in the liver. *Molecules.* 2008;13: 2249–2259.

27. Sakr SA, Mahran HA, Lamfon HA. Protective effect of ginger (*Zingiber officinale*) on adriamycin-induced hepatotoxicity in albino rats. J Med Plant Res. 2011;5(1):133–140.
28. Garouiel M, Fetoui H, Makni FA, Boudawara T, Zeghal N. Cobalt chloride induces hepatotoxicity in adult rats and their suckling pups. Exp Toxicol Pathol. 2011;63(1–2):9–15.
29. Giridharan Bupesh, Chinnaiah Amutha, Sakthivel Vasanth, Natesan Manoharan, Ramalingam Senthil Raja, Raju Krishnamoorthy, Periyasamy Subramanian. Hepatoprotective activity of Red algae *H. muciformis* against experimentally induced hepatotoxicity in albino rats. Brazilian Journal of Biology and Technology. 2012;89:1509-1607.

© 2018 Venkatachalam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/26236>