



Molecular Docking Studies on Phenolic Constituents of *Anethum graveolens* L. Seed Extracts

Paul Andrei Negru^a≡, Sanda Rodica Bota^bⓄ, Oana Delia Stanasel^b##*, Cristian Felix Blidar^cⓄ and Georgeta Serban^a†*

^a *Pharmaceutical Chemistry Department, Faculty of Medicine and Pharmacy, University of Oradea, Romania.*

^b *Chemistry Department, Faculty of Informatics and Sciences, University of Oradea, Romania.*

^c *Biology Department, Faculty of Informatics and Sciences, University of Oradea, Romania.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors GS and SRB conceived and designed the experiments. All authors analyzed the data. Authors GS and SRB wrote the manuscript. All authors have read and approved the final version of the manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i56B33939

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/78663>

Original Research Article

Received 10 September 2021

Accepted 12 December 2021

Published 13 December 2021

ABSTRACT

Background: There are studies indicating that aqueous or hydroalcoholic dill extracts showed higher antioxidant activity compared to other fractions. Molecular docking studies would be relevant to get information on the mechanism of action of the phenolic constituents of *Anethum graveolens* seed extracts as bioactive compounds.

Methodology: In order to perform the docking studies of antioxidant activity of phenolic constituents of *Anethum graveolens* seed extracts, BIOVIA Discovery Studio and AutoDock Vina software were used.

Results: The orientation of flavonoids within Hck and CYP2C9 binding sites has been shown to be the main reason for their inhibitory potency.

≡ *Doctoral Student;*

Ⓞ *Lecturer PhD.;*

Assoc. Prof. PhD.;

† *Assoc. Prof. Habilitate PhD.;*

**Corresponding author: E-mail: ostanasel@uoradea.ro, getaserban_2000@yahoo.com;*

Conclusion: Molecular docking studies indicate that the compounds identified interact with the target enzymes Hck and CYP2C9 at molecular level through their condensed ring systems and hydroxyl substituents and therefore support the antioxidant capacity of the studied phenolic compounds.

Keywords: Dill seeds; extracts; ligands; molecular targets.

1. INTRODUCTION

Clinical studies have shown that very common chronic age-related diseases have increased oxidative stress. The ability of phytochemicals to lower the level of free radicals is associated with their benefits in the prevention of chronic diseases caused by oxidative stress and it may be associated with their antioxidant activity [1,2]. Natural polyphenols, such as phenolic acids, flavonoids and coumarins, are the most important group of secondary metabolites found in many medicinal plants as well as in the human diet and contribute to the antioxidant properties of plants and foods [1,3].

Synthetic phenolic antioxidants are scarcely accepted due to their ineffectiveness and toxicity [4, 5]. As the risk factors for the development of lethal diseases among population are constantly increasing, there is a global tendency to use natural substances from medicinal herbs and food plants as therapeutic antioxidants [4, 5].

Plants used as aromatic herbs or spices are often underestimated by their lack of metabolic profile, as they are often important sources of bioactive compounds [6]. There are two related species of cultivated dill, the European dill (*A. graveolens*) which is widely distributed throughout the world and the Indian dill (*A. sowa*) cultivated throughout the Indian sub-continent, Malaysia and Japan [7,8]. Since the Egyptian period, dill has been used as a condiment and also for medicinal purposes as an ingredient in pain relievers [7,8]. *A. graveolens* is used in India in the preparation of more than 56 ayurvedic medicines, which are usually prescribed in abdominal discomfort, colic and digestive disorders [8], while in Ethiopia dill is one of the plants used in the traditional treatment of kidney diseases [3]. In Romania, the whole dill plant is used in the traditional cuisine to flavor different foods, and the seeds are used as spices for pickles. Hydroalcoholic and hydroglycerin extracts are recommended for emmenagogue, estrogenic, galactagogue, and female hormone regulator, anti-inflammatory, antispasmodic, diuretic, stomachic and carminative effects. Dill

seed extract showed significant protective gastric activities, having anti-ulcer effects in mice by oral administration of hydrochloric acid and absolute ethanol [9], antinociceptive properties in inflammatory pain [10], antioxidant activities [11], anticonvulsant effect against pentylenetetrazole [12], antimicrobial activity against bacteria, such as *Bacillus subtilis*, *Escherichia coli*, *Helicobacter pylori*, *Salmonella sp.*, *Shigella flexneri*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* or fungi such as *Aspergillus flavus*, *Candida albicans*, *Fusarium graminearum* [7,8,13].

Dill seeds contain up to 5% volatile oil – rich in carvone and limonene as major compounds, flavonoids, coumarins, xanthenes and terpenes [3,8]. Several studies have been done to determine the chemical composition of the essential oil of dill seeds and it has been shown to exert antioxidant activity, suggesting a possible application as a natural preservative for food, as well as a protector of the body against oxidative stress [13,14]. However, essential oils are volatile and difficult to store due to low stability, while extracts are easier to prepare, cheaper and more stable. There are studies indicating that aqueous or hydroalcoholic dill extracts showed higher antioxidant activity compared to other fractions [15,16]. To the best of our knowledge, there are few reports concerning the chemical composition of the alcoholic and hydroalcoholic extracts of *Anethum graveolens* seeds and the data differ from author to author, as different components have been analyzed [5,17,12,18]. Dill seeds are marketed in Romania, but it is not yet known how detailed the plant product has been studied from a chemical point of view, so its value as food/nutraceutical. Moreover, dill seeds have many medicinal properties. Based on these considerations, in a previous paper [19], we reported the total phenolics contents and several phenolic compounds identified in methanolic and hydromethanolic extracts from dill seeds cultivated in Romania. Antioxidant activity, as measured by DPPH's free radical scavenging ability, was also investigated. Protein tyrosine kinases (such as Hck) and cytochrome P-450

proteins are enzymes involved in various cellular processes, including oxidative biotransformation in the body, and their inhibition may relieve some clinical symptoms. Although phenolic compounds are known to have antioxidant properties, the exact mechanism by which they act is often not fully understood, and molecular docking studies can provide valuable information in this regard. Therefore, molecular docking studies were performed to obtain some information in an attempt to understand the mechanism of antioxidant activity of dill seed extracts.

2. MOLECULAR DOCKING METHODOLOGY

The crystal structures of Hck (2HCK) and CYP2C9 (1OG5) were obtained from the Research Collaboratory for Structural Bioinformatics Protein Data Bank (RCSB PDB) [20,21] as PDB files and used after removal of all bound water molecules, ligands and cofactors by using BIOVIA Discovery Studio software [22] (BIOVIA, San Diego, CA, USA). Polar hydrogen atoms and Gasteiger charges were added using AutoDock Vina software (Scripps Research, San Diego, CA, USA). All other settings were left as default [23]. The ligands were obtained from the PubChem database in SDF (structure-data file) format and were converted to the PDB format using <https://cactus.nci.nih.gov/translate/>. The ligands were prepared by adding Gasteiger charges, removing non-polar hydrogens and defining all rotatable bonds. Docking was performed by placing the ligands in the binding pocket of the protein (together in one PDB file) and the binding affinity was calculated with AutoDock Vina software. The coordinates of the grid center for 2HCK are 31.47x, 45.34y and 98.79z, grid size 24x, 24y, 24z while the coordinates for 1OG5 are -20.256x, 86.991y and 38.581z, grid size 24x, 24y, 24z. BIOVIA Discovery Studio software (BIOVIA, San Diego, CA, USA) was used to design the 2D diagrams and to analyze the data obtained.

3. RESULTS AND DISCUSSION

Molecular docking is a commonly used method to predict the preferred orientation and binding energy of a small molecule (ligand) to a target macromolecule (receptor). This can be useful for understanding the mechanism of action and the nature of ligand binding to different types of receptors [24]. Molecular docking studies were performed to evaluate the antioxidant potential of the phenolic compounds identified in the

methanolic and hydromethanolic extracts of *A. graveolens* seeds that we have previously studied [19]. The aim of these studies was to establish the relationship between in vitro results and in silico experiments. Two proteins were selected as targets for antioxidant compounds, tyrosine kinase Hck [25] and cytochrome P450, CYP2C9 [26,27].

Protein kinases are enzymes commonly found in eukaryotes that are involved in various cellular processes such as cell cycle, proliferation, metabolism and apoptosis. Activation of protein-tyrosine kinases can induce controlled production of endogenous hydrogen peroxide (H₂O₂) [28]. In contrast, excessive generation of H₂O₂ and other reactive oxygen species (ROS) can alter some enzymes, including protein-tyrosine kinases. The generation of abnormal protein-tyrosine kinases and exacerbation of oxidative stress can lead to damage of biomolecules and pathologies such as cancer, diabetes, neurodegenerative and cardiovascular diseases. Therefore, it can be anticipated that inhibitors that block tyrosine kinases activity and the signaling pathways they activate could alleviate clinical symptoms and provide a useful basis for the development of new drugs [28,29]. Cytochrome P-450 proteins (CYP450) are ubiquitous hemoxygenases that play a critical role in the oxidative biotransformation of a wide range of physiologically important substances as well as exogenous compounds such as drugs, environmental compounds and pollutants [26,27]. CYP2C9 is the predominant isoform of CYP450 with a major contribution to human drug metabolism [27]. CYP450 is also known to generate ROS during arachidonic acid metabolism [30].

Therefore, we performed an in silico evaluation of the binding poses and functional group interactions between the phenolic derivatives identified in *A. graveolens* extracts and the catalytic sites of Hck and CYP2C9 in order to examine the potential molecular interactions between these compounds and the target enzymes. The molecular docking was first conducted on compounds known as reference ligands and the results were compared with these known molecules used as positive controls. Quercetin was chosen as reference ligand for Hck (quercetin-Hck complex, PDB code: 2HCK) [25], while S-warfarin was chosen as the reference ligand for the protein CYP2C9 (S warfarin - CYP2C9 complex, PDB code: 1OG5) [26]. To validate the method, a ligand

model was built and docked into the crystal structure of the target enzyme. The docked ligand model was superimposed on the reference model of ligand taken from the protein data bank. Root mean square deviation (RMSD) between the co-crystallized ligand and the docked ligand showed a value of 1.097 Å for 2HCK (Fig. 1) and 0.89 Å for 1OG5 (Fig. 2), respectively, which fits within the generally accepted values of 1.5-2 Å [30].

The identified phenolic compounds (Fig. 3) were docked within the active site of the target macromolecules and the binding affinity of the different conformations of the inhibitors was calculated with AutoDock Vina software. As can be seen in Table 1, the best docking scores were obtained by compounds with condensed ring systems: quercetin, kaempferol and (±) - catechin belonging to the flavonoid family, followed by the coumarin umbelliferone.




Fig. 1. Data obtained in the validation of molecular docking methods for Hck target enzyme. Quercetin in 2HCK complex, Quercetin binding affinity of - 9.4 kcal/mol, RMSD = 1.097 Å




Fig. 2. Data obtained in the validation of molecular docking methods for CYP2C9 target enzyme. S-warfarin in 1OG5 complex, S-warfarin binding affinity of - 9.9 kcal/mol, RMSD = 0.890 Å




Fig. 3. Phenolic compounds screened and identified in *A. graveolens* seed extracts

The binding interactions of the docked conformations of ligands with target proteins are identified in 2D diagrams obtained using BIOVIA Discovery Studio software (BIOVIA, San Diego, CA, USA). The amino acids at the active site of the target protein as well as the functional groups through which the studied compounds bind to these amino acids can be seen in these diagrams.

Table 1. Binding affinity within 2HCK and 1OG5 of the phenolic compounds identified in *A. graveolens* extracts

Compound	Binding affinity (kcal/mol)	
	2HCK	1OG5
Quercetin	-9.4	-8.7
Kaempferol	-9.0	-8.6
Caffeic acid	-6.4	-6.1
Sinapic acid	-6.8	-6.2
Gallic acid	-5.8	-5.9
Vanillic acid	-6.0	-5.5
(±)-Catechin	-8.7	-8.1
Umbelliferone	-7.1	-6.3
S-warfarin	-	-9.9

The study of 2HCK complex shows that the active site of Hck is coated by LEU273, GLY274, ALA275, VAL281, ALA293, VAL323, THR338, GLU339, PHE340, MET341, GLY344, SER345, ALA390, LEU393 and ALA403 residues. Specific interactions occur between these residues and the condensed rings of quercetin: hydrogen

bonds (GLU339, MET341), Pi-sigma interactions (LEU393, VAL281), Pi-alkyl interactions (ALA293, LEU273, VAL281). In addition, the phenyl ring in position 2 is involved in hydrogen bonds (ALA390), as well as Pi-sigma interactions (VAL281). Several van der Waals interactions occur between the quercetin molecule and the enzyme residues (PHE340, SER345, GLY274, ALA275, ALA403, THR338, VAL323), as well as one carbon-hydrogen bond (GLY344). Due to its similar structure, kaempferol showed a similar binding affinity to the protein (- 9.0 kcal/mol). Comparing the 2D diagrams of these two compounds, it can be seen that the protein binding mode is very similar, suggesting that these flavonoids may inhibit Hck activity with similar potency. The small difference in binding affinity may be caused by the lack of one hydroxyl group on the phenyl ring of kaempferol resulting in the lack of the hydrogen bond with ALA390. Catechin interacted with seven residues common to quercetin (GLU339, THR338, LEU273, ALA403, LEU393, VAL281, GLY344), that are involved in hydrogen bonds, Pi-sigma and Pi-alkyl interactions and therefore revealing a high affinity of this compound for Hck (- 8.7 kcal/mol) (Fig. 4). The binding affinities of the compounds and the control molecule are similar due to the related chemical structures. However, the lower binding affinity of catechin may suggest that the orientation of flavonoids within the catalytic site may influence their inhibitory capacity.




Fig. 4. Interactions of the flavonoids identified in *A. graveolens* extracts with Hck enzyme

Several studies have reported that flavonoids are effective inhibitors of protein kinases [31,32,33] and through this action flavonoids can regulate processes such as cell proliferation, immune response, inflammation or blood clotting, preventing pathologies such as cancer, diabetes, neurodegenerative and cardiovascular diseases [32]. From the 2D diagrams of flavonoid-Hck complexes it can be concluded that the condensed ring systems and their hydroxyl substitutions are important structural features for Hck binding.

The study of the 1OG5 complex shows that the active site of CYP2C9 is surrounded by ARG97, GLY98, ILE99, PHE100, LEU102, ALA103, VAL113, PHE114, LEU208, ILE213, GLN214, ASN217, THR364, SER365, LEU366, PRO367, LEU388 and PHE476 residues which interact with the control ligand S-warfarin through hydrogen bonds (PHE100, ASN217, GLN214), Pi-Pi (PHE114, PHE476) and Pi-alkyl

interactions (PRO367, ARG97, LEU366, ALA103), one carbon - hydrogen bond (ILE99) and van der Waals interactions (GLY98, LEU388, THR364, LEU208, SER365, ILE213, LEU102, VAL113) which is in agreement with published data [26]. Quercetin docked in a different orientation to that of warfarin, with the condensed rings positioned adjacent to PHE476, GLN214, ASN217, ALA477 and LEU208 residues, three of which being involved in hydrogen bonds (PHE476, GLN214 and ASN217). From the list of residues, six of them interact with quercetin (ALA103, PHE114, GLN214, ASN217, PRO367 and PHE476) which indicates a good antioxidant ability of this compound (binding affinity value - 8.7 kcal/mol). Kaempferol docked in an orientation similar to quercetin, making hydrogen bonds with GLN214, ASN217 and VAL113 residues, suggesting that these flavonoids may inhibit CYP2C9 activity with similar potency (binding affinity value - 8.6 kcal/mol), while catechin docked in a more

warfarin-like orientation, forming bonds with common residues in CYP2C9 binding site. Although catechin docked with a similar number of hydrogen bonds (ASN217, PHE476 and LEU208) as warfarin, the binding affinity of catechin (- 8.1 kcal/mol) is lower to that of warfarin (- 9.9 kcal/mol). The unfavorable binding to PHE100 could be a reason. In addition, although the hydroxyl group on the non-planar catechin ring is involved in hydrogen bonds, it is probably unfavorable from an energy point of view (Fig. 5).

These results demonstrate that hydrogen bonds and hydrophobic interactions occur between ligands and target proteins, stabilizing the ligands at the catalytic site and providing high binding affinities. The unique stability of the ligands in the binding sites can be attributed to the large number of Pi-interactions such as Pi-sigma, Pi-anion, Pi-Pi and Pi-alkyl. It also appears that orientation of the flavonoids within Hck and CYP2C9 binding sites, rather than the number of hydrogen bonds formed, may determine their inhibitory potency.




Fig. 5. Interactions of the flavonoids identified in *A. graveolens* extracts with CYP2C9 enzyme

4. CONCLUSIONS

Molecular docking studies provide information on possible molecular targets and support the antioxidant capacity of the studied phenolic compounds, and subsequent chemical analysis of dill seeds will make a valuable contribution to understanding the phytochemical composition and pharmacological properties of this edible plant.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

This research did not receive any specific grants from funding agencies in the public, commercial, or not-for-profit sectors.

COMPETING INTERESTS

The study highlights the efficacy of "*Anethum graveolens*" which is an ancient tradition, used in many countries, including Romania. This ancient concept should be carefully evaluated in the light of modern medical science and may be recommended if found suitable.

REFERENCES

1. Zhang YJ, Gan RY, Li S, Zhou Y, Li AN, Xu DP et al. Antioxidant phytochemicals for the prevention and treatment of chronic diseases. *Molecules* 2015;20(12):21138-21156.
DOI: 10.3390/molecules201219753
2. Forni C, Facchiano F, Bartoli M, Pieretti S, Facchiano A, D'Arcangelo D et al. Beneficial role of phytochemicals on oxidative stress and age-related diseases. *BioMed Research International*. 2019; Article ID 8748253:16.
DOI: 10.1155/2019/8748253
3. Memariani Z, Gorji N, Moeini R, Farzaei MH. Traditional uses. In: Nabavi SM, Sutar I, Barreca D, Khan H (editors). *Phytonutrients in Food: From Traditional to Rationale Usage*. Duxford, UK: Woodhead Publishing, Elsevier, 2020;23-66.
4. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy Reviews* 2010;4(8):118-126.
DOI: 10.4103/0973-7847.70902
5. Tanruean K, Kaewnarin K, Rakariyatham N. Antibacterial and antioxidant activities of *Anethum graveolens* L. dried fruit extracts. *Chiang Mai Journal of Science*. 2014; 41(3):649-660.
6. Fernandes F, Andrade PB, Ferreres F, Gil-Izquierdo A, Sousa-Pinto I, Valentao P et al. The chemical composition on fingerprint of *Glandora diffusa* and its biological properties. *Arabian Journal of Chemistry* 2017;10(5):583-595.
DOI: 10.1016/j.arabjc.2015.01.012
7. Gupta R, Anwer MM, Sharma YK. Dill. In: Peter KV (editor). *Handbook of Herbs and Spices*. Second edition, Volume 1. Cambridge, UK: Woodhead Publishing Ltd. 2012;275-285.
8. Jana S, Shekhawat GS. *Anethum graveolens*: An Indian traditional medicinal herb and spice. *Pharmacognosy Reviews*. 2010;4(8):179-184.
DOI: 10.4103/0973-7847.70915
9. Hosseinzadeh H, Karimi GR, Ameri M. Effects of *Anethum graveolens* L. seed extracts on experimental gastric irritation models in mice. *BMC Pharmacology*. 2002;2(1):21.
DOI: 10.1186/1471-2210-2-21
10. Rezaee-Asl M, Bakhtiarian A, Nikoui V, Sabour M, Ostadhadi S, Yadavar-Nikraves MS et al. Antinociceptive properties of hydroalcoholic extracts of *Anethum graveolens* L. (dill) seed and aerial parts in mice. *Journal of Clinical and Experimental Pharmacology*. 2013;3: 122.
DOI: 10.4172/2161-1459.1000122
11. Koppula S, Choi DK. *Anethum graveolens* L. (Umbelliferae) extract attenuates stress-induced urinary biochemical changes and improves cognition in scopolamine induced amnesic rats. *Tropical Journal of Pharmaceutical Research*. 2011;10(1):47-54.
12. Rostampour M, Ghaffari A, Salehi P, Saadat F. Effects of hydro-alcoholic extract of *Anethum graveolens* seed on pentylentetrazol-induced seizure in adult male mice. *Basic and Clinical Neuroscience* 2014;5 (3):199-204.
13. Stanojević LP, Stankovic MZ, Cvetkovic DJ, Danilovic BR, Stanojević JS. Dill

- (*Anethum graveolens* L.) seeds essential oil as a potential natural antioxidant and antimicrobial agent. *Biologica Nyssana* 2016;7(1):31-39.
DOI: 10.5281/zenodo.159101
14. Rădulescu V, Popescu ML, Ilieș DC. Chemical composition of the volatile oil from different plant parts of *Anethum graveolens* L. (Umbelliferae) cultivated in Romania. *Farmacia*. 2010;58(5):594-600.
 15. Shyu YS, Lin JT, Chang YT, Chiang CJ, Yang DJ. Evaluation of antioxidant ability of ethanolic extract from dill (*Anethum graveolens* L.) flower. *Food Chemistry* 2009;115(2):515-521.
DOI: 10.1016/j.foodchem.2008.12.039
 16. Abbasi Oshaghi E, Khodadadi I, Tavilani H, Goodarzi MT. Aqueous extract of *Anethum graveolens* L. has potential antioxidant and antiglycation effects. *Iranian Journal of Medical Sciences*. 2016; 41(4):328-333.
 17. Jurca Paven CS, Radu D, Alexa E, Pintilie S, Ravis A. *Anethum graveolens* – an important source of antioxidant compounds for food industry. In: 18th International Multidisciplinary Scientific GeoConference SGEM Proceedings – Advances in Biotechnology; Albena, Bulgaria. 2018;11-18.
 18. Ishikawa T, Kudo M, Kitajima J. Water-soluble constituents of dill. *Chemical and Pharmaceutical Bulletin* 2002;50(4):501-507.
DOI: 10.1248/cpb.50.501
 19. Bota SR, Stanasel OD, Blidar CF, Serban G. Phenolic constituents of *Anethum graveolens* seed extracts: chemical profile and antioxidant effect studies. *Journal of Pharmaceutical Research International* 2021;33(54B):168-179.
DOI: 10.9734/JPRI/2021/v33i54B33777
 20. Protein Data Bank. 2HCK. Src family kinase Hck-quercetin complex [online]. Available: <https://www.rcsb.org/structure/2HCK>
(Accessed 22 October 2020)
 21. Protein Data Bank. 1OG5. Structure of human cytochrome P450 CYP2C9 [online]. Available: <https://www.rcsb.org/structure/1OG5>
(Accessed 22 October 2020)
 22. Dassault Systèmes BIOVIA, Discovery Studio v19.1.0.18287. Dassault Systemes: San Diego, CA, USA; 2020
 23. Trott O, Olson AJ. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization and multithreading. *Journal of Computational Chemistry*. 2010; 31(2):455-461.
DOI: 10.1002/jcc.21334
 24. Ferreira LG, dos Santos RN, Oliva G, Andricopulo AD. Molecular docking and structure-based drug design strategies. *Molecules*. 2015;20(7):13384-13421.
DOI: 10.3390/molecules200713384
 25. Sicheri F, Moarefi I, Kuriyan J. Crystal structure of the Src family tyrosine kinase Hck. *Nature*. 1997;385(6617):602-609.
DOI: 10.1038/385602a0
 26. Williams PA, Cosme J, Ward A, Angove HC, Matak Vinković D, Jhoti H. Crystal structure of human cytochrome P450 2C9 with bound warfarin. *Nature*. 2003; 424(6947):464-468.
DOI: 10.1038/nature01862
 27. Tarcsay A, Kiss R, Keseru G. Site of metabolism prediction on cytochrome P450C9: A knowledge-based docking approach. *Journal of Computed-Aided Molecular Design*. 2010;24(5):399-408.
DOI: 10.1007/s10822-010-9347-3.
 28. Truong TH, Carroll KS. Redox regulation of protein kinases. *Critical Reviews in Biochemistry and Molecular Biology*. 2013; 48(4):332-356.
DOI: 10.3109/10409238.2013.790873
 29. Li J, Pang Q. Oxidative stress-associated protein tyrosine kinases and phosphatases in Fanconi Anemia. *Antioxidants and Redox Signaling*. 2014;20(14):2290-2301.
DOI: 10.1089/ars.2013.5715
 30. Costa JS, Ramos RS, Costa KSL, Brasil DSB, Silva CHTP, Ferreira EFB et al. An in silico study of the antioxidant ability for two caffeine analogs using molecular docking and quantum chemical methods. *Molecules*. 2018;23(11):2801-2817.
DOI: 10.3390/molecules23112801
 31. Hou DX, Kumamoto T. Flavonoids as protein kinase inhibitors for cancer chemoprevention: Direct binding and molecular modeling. *Antioxidants and Redox Signaling*. 2010;13(5):691-719.
DOI: 10.1089/ars.2009.2816.

32. Wright B, Watson KA, McGuffin LJ, Lovegrove JA, Gibbins JM. GRID and docking analyses reveal a molecular basis for flavonoid inhibition of Src family kinase activity. *Journal of Nutritional Biochemistry*. 2015;26(11):1156-1165.
DOI: 10.1016/j.jnutbio.2015.05.004
33. Baier A, Nazaruk J, Galicka A, Szyszka R. Inhibitory influence of natural flavonoids on human protein kinase CK2 isoforms: effect of the regulatory subunit. *Molecular and Cellular Biochemistry*. 2018;444(1):35-42.
DOI: 10.1007/s11010-017-3228-1

© 2021 Negru et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78663>