



Evaluating the Effect of Seed Priming Treatments on Yield and Quality of Wheat Seed (*Triticum aestivum* L.) under Rainfed Conditions

Shravan Kumar ^{a++*}, A. L. Jatav ^{a#}, C. L. Maurya ^{a#},
Ravi Dixit ^{a++}, Shailendra Pratap Singh ^{b++}, Bhayankar ^{c++},
Jayant ^{d++}, Sanjay Babu ^{a++}, Deepu ^{c++}
and Dharendra Kumar ^{e++}

^a Department of Seed Science and Technology, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur- 208002, India.

^b Department of Soil Conservation & Water Management, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur- 208002, India.

^c Department of Agronomy, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur- 208002, India.

^d Department of Genetics & Plant Breeding, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur- 208002, India.

^e Department of Plant Physiology, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur- 208002, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i102926>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124242>

Original Research Article

Received: 24/07/2024

Accepted: 28/09/2024

Published: 28/09/2024

⁺⁺ Research Scholar;

[#] Professor;

*Corresponding author: E-mail: skcsa9026@gmail.com;

Cite as: Kumar, Shravan, A. L. Jatav, C. L. Maurya, Ravi Dixit, Shailendra Pratap Singh, Bhayankar, Jayant, Sanjay Babu, Deepu, and Dharendra Kumar. 2024. "Evaluating the Effect of Seed Priming Treatments on Yield and Quality of Wheat Seed (*Triticum Aestivum* L.) under Rainfed Conditions". *Journal of Experimental Agriculture International* 46 (10):73-80. <https://doi.org/10.9734/jeai/2024/v46i102926>.

ABSTRACT

The present study was conducted on wheat variety "K-1317" and was procured from Student instructional Farm, C.S. Azad University of Agriculture & Technology, Kanpur. The experiment was conducted in Rabi 2022-23 and 2023-24. The seeds were used for pre-sowing seed treatments, T₀ Control (Un soaking), T₁ Soaking with tap water for 12hrs, T₂ Soaking with tap water for 24hrs, T₃ Soaking with *Bacillus Subtilis* for 12hrs, T₄ Soaking with *Bacillus subtilis* for 24hrs, T₅ Soaking with *Azotobacter* for 12hrs, T₆ Soaking with *Azotobacter* for 24hrs, T₇ Soaking with NaCl @1% for 12hrs, T₈ Soaking NaCl @1% for 24hrs, T₉ Soaking with ZnSo₄ @1% for 12hrs, T₁₀ Soaking with ZnSo₄ @1% for 24hrs, T₁₁ Soaking with GA₃ @(50ppm) for 12hrs, T₁₂ Soaking with GA₃ @(50ppm) for 24hrs & T₁₃ Soaking Auxin (IAA) @(50ppm) for 12hrs & T₁₄ Soaking IAA @(50ppm) for 24hrs. Pre-sowing seed treatment or priming was done by soaking of required quantity of seeds of wheat variety K- 1317 in tap water, biologicals, chemicals, and hormonal treatments in concentration for 12hrs and 24hrs in ratio of 1:1 (Kg of seeds/volume of solution) by using wet gunny bags. Then the treated or primed (soaked) seeds were dried in shade to maintain the seed moisture content approximately 12 or 13%. Seed dressing was done on primed and untreated (control) by Thiram (2.5%). Seeds priming was GA₃@ (50ppm) for 12hrs on wheat variety K-1317 significantly present results the yield and quality of seed. The best results in found under the treatment T₁₁ Soaking with GA₃ @(50ppm) for 12hrs, in productive tillers, number seeds/spike, number of spikelet's/spikes, spike length, grain yield and harvest index, protein content in 2022-2023. and the at par treatments T₁₀, T₁₂, T₁₃ and T₁₄.

Keywords: *Wheat crop; Auxin; GA3; Bacillus subtilis.*

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is a self-pollinated crop belonging to the *Poaceae* family and one of the most leading cereals of many countries of the world including India. It has been described as the 'King of cereals' because of the acreage it occupies, high productivity and the prominent position it holds in the international food grain trade. It is the most important food crop of India and is a main source of protein and energy. According to the earliest historic records, wheat was an important cultivated cereal in South-Western Asia, its geographical center of origin. The central Asia, Near East, Mediterranean and Ethiopian regions are the world's most important centers of diversity of wheat and its related species. Hindukush area is the center of diversity of hexaploid wheat [1].

In India, during 2020-21 production of wheat was recorded to 109.52mt with average productivity of 35.00q/ha from an area of 31.61 million ha which constitute 36 percent of the country's total food grain production 308.65mt [2]. In Uttar Pradesh, the total wheat production was 35.50mt and average productivity was 32.42q/ha., with the area of 9.78 million ha. The perusal of state wise production indicates that Uttar Pradesh tops the list with 35.50 mt, followed by Madhya Pradesh (17.62 mt), Punjab (17.14mt). These top three states together contributed about 64 per cent of the total production [3].

Micronutrient deficiency is also known as Hidden Hunger, is one of the most important challenges facing humanity today [4]. or reported the seed priming with GA₃ enhance emergence, stand establishment, tillering, allometry, grain and straw yields, and harvest index by Assefa et al. [5]. Seed priming showed positive response to different priming treatments recorded significantly higher pooled average field emergence (96.42%), number of effective tillers (7.75), spike length (12.99 cm), seed yield per plant (13.55 g), seed yield per plot (2.45 kg) during both the years followed by GA₃ (50 ppm) primed seeds which recorded at par values for almost all the seed yield attributes by Avinash karjule et al. [6].

The wheat grown in India is spring type belong to species *Triticum aestivum* (bread wheat). Unlike other cereals, wheat contains a high amount of gluten, the protein that provides the elasticity necessary for excellent bread making. It has good nutrition profile with 12.1 percent protein, 1.8 per cent lipids, 1.8 percent ash, 2.0 per cent reducing sugars, 6.7 percent pentose's, 59.2 percent starch, 70 percent total carbohydrates and provides 314KCal/100g of food. It is also a good source of minerals and vitamins viz., calcium (37 mg/100g), iron (4.1 mg/100g), thiamine (0.45mg/100g), riboflavin (0.13mg/100g) and nicotinic acid (5.4mg/100mg) (Lorenz and Kulp, 1991). Hard wheat had high protein (10-17%) and yields flour rich gluten, making it particularly suitable for yeast breads.

The low-protein (6 to 10%) softer type yields flour lower in gluten and therefore, suited better for tender baked products, such as biscuits, pastries and cakes.

Priming levels of Zn (0% ZnSO₄-water soaking, 1% ZnSO₄, 2% ZnSO₄ and 3% ZnSO₄) and three priming durations (4 hours, 8 hours and 12 hours) were compared with basal dose of recommended NPK + ZnSO₄ and farmers' practice (absolute control). The yield of green peas, maize equivalent yield was maximum due to priming with 1% ZnSO₄ for 12 hours duration, however, yield of maize grains was highest with treatment combination of 2% ZnSO₄ priming for 12 hours duration by Munish Sharma and D. K. Parmar [7].

Seed priming is one of the advanced methods for accelerating the germination by which, it is possible to increase the ability of seeds to germinate, grow and enhance to yield under moisture stress condition. Farmer's faces moisture stress during early plant growth stages which were resulted in less plant population and the Seed priming is commonly used to reduce the time between seed sowing and seed ling emergence and to synchronize seed ling emergence. The rationale is that sowing the soaked seed de creases the time needed for germination and may al low the seed lings to escape from the deteriorating soil physical conditions. Keeping in view the above facts, the experiment is planned to carry out with the following objectives: -

- ❖ To find out the suitable seed priming treatment for assuring better field emergence.
- ❖ To study the effect of seed priming treatment on plant growth related parameters.
- ❖ To assess the effect of seed priming treatment for getting higher yield and quality attributing parameters.
- ❖ To see the effect of priming treatments on seed quality of wheat.
- ❖ Effect of seed priming treatments on seed health.

2. MATERIALS AND METHODS

The present study was conducted on wheat variety "K-1317" and was procured from Student instructional Farm, C.S. Azad University of Agriculture & Technology, Kanpur. The experiment was conducted in Rabi 2022-23 and 2023-24. The semi-arid climate and rich alluvial

soil characterize this northern zone. About 935 mm of rain falls on the region each year on average. Relative humidity (7 am) is relatively constant at about 80-90% from July to the end of March, gradually declines to about 40-50% by the end of April, and remains at 80% until June, even though temperatures in May and June can reach 44°C to 47°C or higher. The seeds were used for pre-sowing seed treatments, T₀ Control (Un soaking), T₁ Soaking with tap water for 12hrs, T₂ Soaking with tap water for 24hrs, T₃ Soaking with *Bacillus Subtilis* for 12hrs, T₄ Soaking with *Bacillus subtilis* for 24hrs, T₅ Soaking with *Azotobacter* for 12hrs, T₆ Soaking with *Azotobacter* for 24hrs, T₇ Soaking with NaCl @1% for 12hrs, T₈ Soaking NaCl @1% for 24hrs, T₉ Soaking with ZnSo₄ @1% for 12hrs, T₁₀ Soaking with ZnSo₄ @1% for 24hrs, T₁₁ Soaking with GA₃ @(50ppm) for 12hrs, T₁₂ Soaking with GA₃ @(50ppm) for 24hrs & T₁₃ Soaking Auxin (IAA) @(50ppm) for 12hrs & T₁₄ Soaking IAA @(50ppm) for 24hrs. Pre-sowing seed treatment or priming was done by soaking of required quantity of seeds of wheat variety K- 1317 in tap water, biologicals, chemicals, and hormonal's treatments in concentration for 12hrs and 24hrs in ratio of 1:1 (Kg of seeds/volume of solution) by using wet gunny bags. Then the treated or primed (soaked) seeds were dried in shade to maintain the seed moisture content approximately 12 or 13%. Seed dressing was done on primed and untreated (control) by Thiram (2.5%). Primed seeds along with control (untreated) were sown in 25 November in rabi season 2022-23 and 13 November in rabi season 2023-24 under rainfed conditions and apply the FYM @ 12 tonnes, 120kg N, 60kg P₂O₅ and 40 kg K per ha. There were 3 replications by using RBD design for field trail. The crop was raised by using all required agronomical practices. Mature crop was harvested in the last week of April 2023 & 2024. Processed seeds were examined for the quality parameters in three replications at Seed Testing Laboratory of C.S. Azad University of Agriculture & Technology, Kanpur. RBD design was used for statistical analysis. The following field observed recorded, Number of productive tillers/ plot, number of seeds/spike, number of spikelet's/spike, spike length, grain yield kg/hac, harvest index and protein content.

3. RESULTS AND DISCUSSION

3.1 Number of Productive Tillers m⁻¹

Overall mean data presented in Table 1. reveal that all the treatments showed significant differences from T₀ (control). The critical analysis

of data revealed that significant differences were observed among all treatments. Priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs has showed superior (284.33) on number of productive tillers followed by Priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs (283.50) and second followed by (T₁₃) IAA@ (50ppm) soaked for 12hrs (282.83) while the least number of productive tillers was recorded in control (T₀) (273.50). From the analysis, it also perceptible that priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs and priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs increased the number of productive tillers by 10.83 and 10.00 respectively over control. The find similar findings, it can be affirmed that the (*Azotobacter* and PSB) applied seed treatment was recorded in T₃ CG2006 was recorded, which performed well in plant height, spike length, number of seeds per spike, and grain yield, Gayatri et. al. [8] & Avinash karjule et al. [6].

3.2 Number of Seeds Spike⁻¹

Overall mean data presented in Table 1. reveal that all the treatments showed significant differences from T₀ (control). The critical analysis of data revealed that significant differences were observe among all treatments. Priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs has showed superior (44.65) on number of seeds per spike followed by Priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs (44.53) and second followed by (T₁₃) IAA@ (50ppm) soaked for 12hrs (43.72) while the least number of seeds per spike was recorded in control (T₀) (39.38). From the analysis, it also perceptible that priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs and priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs increased the number of productive tillers by 5.27 and 5.19 respectively over control. The find similar findings, it can be affirmed that the (*Azotobacter* and PSB) applied seed treatment was recorded in T₃ CG2006 was recorded, which performed well in plant height, spike length, number of seeds per spike, and grain yield, Gayatri et. al. [8] & Avinash karjule et. al. [6].

3.3 Number of Spikelet's Spike⁻¹

Overall mean data presented in Table 1. reveal that all the treatments showed significant differences from T₀ (control). The critical analysis of data revealed that significant differences were observe among all treatments. Priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs has showed superior (18.82) on number of spikelet's per spike followed by Priming with (T₁₂) GA₃

@ (50ppm) soaked by 24hrs (18.27) and second followed by (T₁₃) IAA@ (50ppm) soaked for 12hrs (117.80) while the least number of spikelet's per spike was recorded in control (T₀) (14.67). From the analysis, it also perceptible that priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs and priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs increased the number of productive tillers by 4.15 and 3.6 respectively over control. The find similar findings, it can be affirmed that (*Azotobacter* and PSB) applied seed treatment was recorded in T₃ CG2006 was recorded, which performed well in plant height, spike length, number of seeds per spike, and grain yield, Gayatri et. al. [8] & Avinash karjule et. al. [6].

3.4 Spike Length (cm)

Overall mean data presented in Table 1. reveal that all the treatments showed significant differences from T₀ (control). The critical analysis of data revealed that significant differences were observe among all treatments. Priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs has showed superior (11.33) on spike length followed by Priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs (11.05) and second followed by (T₁₃) IAA@ (50ppm) soaked for 12hrs (10.13) while the least spike length was recorded in control (T₀) (8.00). From the analysis, it also perceptible that priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs and priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs increased the number of productive tillers by 3.33 and 3.05 respectively over control. The find similar findings, it can be affirmed that (*Azotobacter* and PSB) applied seed treatment was recorded in T₃ CG2006 was recorded, which performed well in plant height, spike length, number of seeds per spike, and grain yield, Gayatri et al. [8] & Avinash karjule et al. [6].

3.5 Grain Yield kg/hac

Overall mean data presented in Table 1 reveal that all the treatments showed significant differences from T₀ (control). The critical analysis of data revealed that significant differences were observe among all treatments. Priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs has showed superior (5104.88) on grain yield kg /hac. followed by Priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs (4998.98) and second followed by (T₁₃) IAA@ (50ppm) soaked for 12hrs (4889.56) while the least grain yield kg/hac. was recorded in control (T₀) (4199.56). From the analysis, it also percentible that priming with (T₁₁)

Table 1. Effect of seed priming treatments on Number of productive tillers per square meter, Number of seeds spike⁻¹ Number of spikelet's per spike and Spike length (cm) in Wheat variety K-1317 under rainfed Condition.

| Treatments | Number of productive tillers per square meter | | | Number of seeds spike ⁻¹ | | | Number of spikelet's per spike | | | Spike length (cm) | | |
|--|---|---------|--------|-------------------------------------|---------|--------|--------------------------------|---------|--------|-------------------|---------|--------|
| | 2022-23 | 2023-24 | Pooled | 2022-23 | 2023-24 | pooled | 2022-23 | 2023-24 | pooled | 2022-23 | 2023-24 | pooled |
| T ₀ Control | 273.33 | 273.67 | 273.50 | 39.21 | 39.55 | 39.38 | 14.137 | 15.21 | 14.67 | 7.90 | 8.10 | 8.00 |
| T ₁ Tap water (12hrs) | 274.00 | 274.33 | 274.17 | 40.28 | 40.98 | 40.63 | 15.40 | 15.54 | 15.47 | 8.47 | 8.70 | 8.58 |
| T ₂ Tap water (24hrs) | 273.33 | 274.00 | 273.67 | 40.12 | 41.01 | 40.57 | 15.31 | 15.49 | 15.40 | 8.76 | 9.12 | 8.94 |
| T ₃ Bacillus subtilis (12hrs) | 278.23 | 279.50 | 278.87 | 41.14 | 41.65 | 41.40 | 16.39 | 16.78 | 16.59 | 8.83 | 9.27 | 9.05 |
| T ₄ Bacillus subtilis (24hrs) | 277.86 | 278.98 | 278.42 | 40.65 | 41.35 | 41.00 | 15.80 | 16.88 | 16.34 | 8.40 | 8.63 | 8.52 |
| T ₅ Azotobacter (12hrs) | 276.33 | 276.67 | 276.50 | 40.98 | 41.00 | 40.99 | 15.99 | 16.08 | 16.04 | 9.09 | 9.74 | 9.42 |
| T ₆ Azotobacter (24hrs) | 275.67 | 276.00 | 275.83 | 40.39 | 41.00 | 40.70 | 16.11 | 16.58 | 16.35 | 8.50 | 7.63 | 8.07 |
| T ₇ NaCl2 @2.5% (12hrs) | 276.33 | 275.96 | 276.15 | 40.26 | 40.59 | 40.42 | 15.61 | 15.78 | 15.70 | 8.57 | 8.60 | 8.58 |
| T ₈ NaCl2 @2.5% (24hrs) | 276.00 | 275.33 | 275.67 | 40.01 | 40.45 | 40.23 | 15.57 | 15.69 | 15.63 | 8.85 | 9.30 | 9.08 |
| T ₉ ZnSo4 @2.5%(12hrs) | 282.33 | 282.67 | 282.50 | 42.23 | 42.86 | 42.55 | 16.71 | 17.21 | 16.96 | 9.53 | 9.87 | 9.70 |
| T ₁₀ ZnSo4 2.5% (24hrs) | 282.00 | 282.33 | 282.17 | 42.03 | 43.00 | 42.52 | 16.50 | 15.60 | 16.05 | 8.98 | 9.07 | 9.02 |
| T ₁₁ GA3 @50ppm (2hrs) | 284.00 | 284.67 | 284.33 | 44.17 | 45.12 | 44.65 | 18.50 | 19.13 | 18.82 | 11.03 | 11.62 | 11.33 |
| T ₁₂ GA3 @50ppm (24hrs) | 283.33 | 283.67 | 283.50 | 44.07 | 44.98 | 44.53 | 18.03 | 18.50 | 18.27 | 10.89 | 10.63 | 10.76 |
| T ₁₃ IAA @50ppm (12hrs) | 282.67 | 283.00 | 282.83 | 43.54 | 43.89 | 43.72 | 17.37 | 18.23 | 17.80 | 9.70 | 10.03 | 9.87 |
| T ₁₄ IAA @50ppm (24hrs) | 281.33 | 281.67 | 281.50 | 43.38 | 43.01 | 43.20 | 16.60 | 17.23 | 16.92 | 8.94 | 8.90 | 8.92 |
| C.D. | 7.68 | 7.56 | 7.62 | 2.09 | 2.101 | 2.10 | 1.10 | 1.07 | 1.08 | 0.99 | 0.79 | 0.89 |
| C.V. | 1.64 | 1.62 | 1.63 | 3.00 | 2.948 | 2.97 | 4.01 | 3.82 | 3.92 | 6.50 | 5.06 | 5.78 |

Table 2. Effect of seed priming treatments on Grain Yield, Harvest index and Protein content in Wheat variety K-1317 under rainfed condition

| Treatments | Grain Yield kg /hac. | | | Harvest index | | | Protein content (%) | | |
|--|----------------------|----------|---------|---------------|---------|--------|---------------------|---------|--------|
| | 2022-23 | 2023-24 | pooled | 2022-23 | 2023-24 | pooled | 2022-23 | 2023-24 | pooled |
| T ₀ Control | 4,142.89 | 4,199.56 | 4171.22 | 35.05 | 35.88 | 35.47 | 10.20 | 10.29 | 10.25 |
| T ₁ Tap water (12hrs) | 4,309.00 | 4,309.21 | 4309.11 | 36.30 | 36.50 | 36.40 | 10.34 | 10.39 | 10.37 |
| T ₂ Tap water (24hrs) | 4,221.01 | 4,148.98 | 4184.99 | 35.98 | 36.20 | 36.09 | 10.09 | 10.20 | 10.14 |
| T ₃ Bacillus subtilis (12hrs) | 4,685.40 | 4,756.26 | 4720.83 | 41.87 | 42.57 | 42.22 | 11.32 | 11.57 | 11.44 |
| T ₄ Bacillus subtilis (24hrs) | 4,659.68 | 4,731.11 | 4695.39 | 40.15 | 40.89 | 40.52 | 11.32 | 11.52 | 11.42 |
| T ₅ Azotobacter (12hrs) | 4,498.56 | 4,593.25 | 4545.90 | 38.97 | 39.57 | 39.27 | 10.68 | 10.74 | 10.71 |
| T ₆ Azotobacter (24hrs) | 4,435.56 | 4,489.25 | 4462.41 | 38.57 | 38.97 | 38.77 | 10.30 | 10.54 | 10.42 |
| T ₇ NaCl2 @2.5% (12hrs) | 4,375.40 | 4,456.89 | 4416.14 | 36.99 | 37.58 | 37.28 | 10.19 | 10.38 | 10.28 |
| T ₈ NaCl2 @2.5% (24hrs) | 4,286.85 | 4,302.79 | 4294.82 | 36.76 | 36.98 | 36.87 | 10.37 | 10.35 | 10.36 |
| T ₉ ZnSo4 @2.5%(12hrs) | 4,861.82 | 4,856.56 | 4859.19 | 41.98 | 42.35 | 42.16 | 10.45 | 10.67 | 10.56 |
| T ₁₀ ZnSo4 2.5% (24hrs) | 4,742.56 | 4,800.23 | 4771.40 | 40.79 | 41.86 | 41.33 | 10.32 | 10.51 | 10.42 |
| T ₁₁ GA3 @50ppm (2hrs) | 5,077.16 | 5,104.88 | 5091.02 | 45.89 | 46.59 | 46.24 | 11.33 | 11.41 | 11.37 |
| T ₁₂ GA3 @50ppm (24hrs) | 4,941.16 | 4,998.98 | 4970.07 | 44.57 | 45.02 | 44.80 | 11.30 | 11.37 | 11.33 |
| T ₁₃ IAA @50ppm (12hrs) | 4,849.56 | 4,889.56 | 4869.56 | 42.88 | 43.21 | 43.05 | 10.35 | 10.53 | 10.44 |
| T ₁₄ IAA @50ppm (24hrs) | 4,805.25 | 4,835.27 | 4820.26 | 43.01 | 43.50 | 43.26 | 10.21 | 10.40 | 10.30 |
| C.D. | 288.526 | 256.403 | 272.46 | 2.76 | 2.44 | 2.60 | 0.44 | 0.36 | 0.40 |
| C.V. | 3.737 | 3.293 | 3.52 | 4.10 | 3.59 | 3.85 | 2.89 | 2.33 | 2.61 |

GA₃ @ (50ppm) soaked for 12hrs and priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs increased the number of productive tillers by 9.05 and 7.99 respectively over control. The find similar findings, it can be affirmed that the seed priming for 8 h provided higher seed yield by 12.0% in the first year and by 5.9% in the second year compared with non-primed control. Christos A. D. et al. [9] and (*Azotobacter* and PSB) applied seed treatment was recorded in T₃ CG2006 was recorded, which performed well in plant height, spike length, number of seeds per spike, and grain yield, Gayatri et al. [8] & Avinash karjule et. al. [6].

3.6 Harvest Index

Overall mean data presented in Table 2. reveal that all the treatments showed significant differences from T₀ (control). The critical analysis of data revealed that significant differences were observe among all treatments. Priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs has showed superior (46.24) on harvest index followed by Priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs (44.80) and second followed by (T₁₄) IAA@ (50ppm) soaked for 24hrs (43.26) while the least harvest index was recorded in control (T₀) (35.47). From the analysis, it also perceptible that priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs and priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs increased the number of productive tillers by 10.77 and 8.79 respectively over control. The find similar findings, it can be affirmed that the and (*Azotobacter* and PSB) applied seed treatment was recorded in T₃ CG2006 was recorded, which performed well in plant height, spike length, number of seeds per spike, and grain yield and harvest index & Avinash Karjule et al. [6].

3.7 Protein Content (%)

Overall mean data presented in Table 2. reveal that all the treatments showed significant differences from T₀ (control). The critical analysis of data revealed that significant differences were observe among all treatments. Priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs has showed superior (42.81) protein content followed by Priming with (T₁₂) GA₃ @ (50ppm) soaked by 24hrs (42.09) and second followed by (T₁₃) Auxin (IAA) @ (50ppm) soaked for 12hrs (41.63) while the least protein content was recorded in control (T₀) (38.06). From the analysis, it also perceptible that priming with (T₁₁) GA₃ @ (50ppm) soaked for 12hrs and priming with (T₁₂) GA₃ @ (50ppm) soaked for 24hrs increased

the number of productive tillers by 4.75 and 4.03 respectively over control [10-14]. The find similar findings, it can be affirmed that the impact of priming, biofertilizers with different nitrogen levels on growth, yield and nutrient uptake of late sown wheat Bishnupriya Patra et al. [15] & Avinash Karjule et. al. [6] and Similar results were also obtained for protein content in grain (12.95 to 13.22 %) and protein yield (632.7 to 678.0 kg ha⁻¹) with *Azotobacter* inoculation by P. K. Jaga et. al. [16].

4. SUMMARY AND CONCLUSION

Seed priming with GA₃ also showed significant improvement over unprimed seeds for the best result found treatment T₁₁ in Number of productive tillers 1m² area, number of seeds/spike, number of spikelet's/spikes, spike length (cm), grain yield kg/hac, harvest index and protein content (%), in 2022 (284.0, 44.27, 18.50, 11.03, 5077.16, 45.89 and 11.33). and 2023 (284.67, 45.12, 19.13, 11.62, 5104.88, 46.59 and 11.41) and pooled (284.33, 44.65, 18.82, 11.33, 5091.02, 46.24 and 11.37) respectively. Whereas seed yield is concerned, GA₃ soaking for 12hrs (50 ppm) primed seed application most suitable to the farmers for harvesting the highest grain yield, yield attributes and for enhancing the value of all quality seed parameters. But the application of GA₃ hormone more costly to farmers for commercial cultivation which is more profitable by increasing the yield and quality of seed and a most common limits to lack of knowledge about these applicable techniques. Significantly positive as well as negative desired traits associated with yield as might be used to improve the yield and seed quality and quality parameters.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kundu S, Nagrajan S. Distinguish characters of Indian wheat varieties. Research bulletin no. 4, Directorate of Wheat research Karnal, India; 1996.

2. Anonymous. Fourth advance estimates of production of food grains for 2021-22 as on 15-05-2022. Director of Economics and Statistics, DAC and FW, Ministry of Agriculture and Farmer Welfare, Govt. of India; 2021-22a.
3. Anonymous. AICRP on NSP Report; 2021-22b. Available: [https://aicrp.icar.gov.in/nsp/enhancement on variety-replacement-rate-vrr-in-major-field-crops/](https://aicrp.icar.gov.in/nsp/enhancement%20on%20variety-replacement-rate-vrr-in-major-field-crops/)
4. White PJ, Broadley MR. Biofortification of crops with seven mineral elements often lacking in human diets iron, zinc, copper, calcium, magnesium, selenium and iodine. *New Phytol.* 2009;182:49-84.
5. Assefa MK, Hunje R, Koti RV. Enhancement of seed quality in soybean following priming treatment. *Karnataka Journal of agriculture Science*, 2010;23:787-89.
6. Karjule A, Kalyanrao Sasidharan N, Patel DA. Effect of different seed priming treatments on yield attributes of wheat (*Triticum aestivum*). *Indian Journal of Agricultural Sciences*; 2019. DOI:10.56093/ijasv.89i10.94597.
7. Sharma M, Parmar DK. Effect of seed priming with zinc sulfate on yield and quality parameters of rainfed maize-pea sequence under mid hill conditions of Himachal Pradesh. *Journal of Pharmacognosy and Phytochemistry, JPP.* 2018;7(1):1401-1407.
8. Gayatri Dr. Singh AK, Verma SON, Sahu U. Effect of *azotobacter* and phosphate solubilizing bacteria on the yield of different wheat (*Triticum aestivum* L.) Cultivar. *The Pharma Innovation Journal.* 2022;11(1):1629-1633.
9. Damalas CA, Koutroubas SD, Sideris Fotiadis S. Hydro-priming effects on seed germination and field performance of Faba Bean in Spring Sowing; 2019
10. Adhikari B, Olorunwa OJ, Barickman CT. Seed Priming Enhances Seed Germination and Morphological Traits of *Lactuca sativa* L. under Salt Stress. 2022;1:74–86. Available: <https://doi.org/10.3390/seeds1020007>.
11. Bassi G, Sharma S, Gill BS. Pre-sowing seed treatment and quality in- vigation in soybean (*Glycine max* L.). *Seed Research.* 2011;31:81-84.
12. Lorenz K, Kulp K. Breakfast cereals, handbook of cereal science and technology. Library of Cogress Newyork. 1991;8247-8358.
13. Singh BA, Gangwar CBS, Singh P, Maurya CL. Effect of seed priming on quality parameters of wheat (*Triticum aestivum* L.) seeds harvested under irrigated & rainfed conditions. *Journal of Pharmacognosy and Phytochemistry.* 2017;6(4):1646-1650.
14. Singh S, Singh Mor V, Kumar ABR, Singh G, Digamber Evaluating the effect of various seed treatment approaches on seed quality of wheat (*Triticum aestivum* L.). *The Pharma Innovation Journal.* 2023;12(10):2439-2444.
15. Patra B, Singh J. Effect of Priming, Biofertilizers and Nitrogen Levels on Yield and Nutrient Uptake by Wheat. *International Journal of Current Microbiology and Applied Sciences.* 2018;7(7). ISSN: 2319-7706 DOI:10.20546/ijcmas.2018.707.168.
16. Jaga PK, Sharma S, Patel Y. Response of wheat (*Triticum aestivum*) to *Azotobacter* inoculation and nitrogen in soils of Vidisha, Madhya Pradesh. *Annals of Plant and Soil Research.* 2017;19(1):42-45.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/124242>