



Ankylosed Pubic Symphysis: A Case Report of an Idiopathic Clinico-radiological Mismatch in an Emergency Trauma

Lavindra Tomar ^a, Gaurav Govil ^{a*} and Pawan Dhawan ^a

^a *Department of Orthopaedics, Max Super Specialty Hospital, 108 A, I.P. Extension, Patparganj, Delhi - 110092, India.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors LT, GG and PD were involved with case management. Author GG collected the data and wrote the first draft. All authors did the literature searches, read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Hab. Mariusz Cycon, Medical University of Silesia, Poland.

Reviewers:

(1) Tom Adejoh, Federal University of Technology, Nigeria.

(2) Dinesh Dhar, Rustaq Hospital, Oman.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here: <https://www.sdiarticle5.com/review-history/81916>

Case Study

Received 25 October 2021

Accepted 27 December 2021

Published 28 December 2021

ABSTRACT

The aim of the case presented was to highlight the assessment of an incidental radiographic finding of fused symphysis for emergency clinicians and trauma surgeons involved in the assessment and management of emergency trauma cases. The objective was to understand the symphyseal affections and to review the literature for approach and management strategy for an ankylosed pubis symphysis. A road traffic accident case of a 62-year old male presented with head, face and pelvic trauma. Standard trauma protocol evaluation was done. Computerised tomography of head revealed basal ganglia bleed and facial tomographic reconstruction was normal. No fractures were evident on the radiography of the pelvis however, an incidental ankylosed symphysis pubis was revealed. Head injury required intensive care management. The incidental radiographic finding had no history suggestive of any possible aetiological cause leading to the ankylosis of pubic symphysis. A clinico-radiological mismatch presented with an idiopathic asymptomatic symphyseal ankylosis. A broader clinician's perspective during the radiological evaluation of an emergency trauma is needed. The case highlights the necessity of taking a detailed and thorough clinical history. The trauma surgeon and the emergency physician need to be aware of the pathological affections of the pubis symphysis leading to ankylosis to strategize for an effective management.

*Corresponding author: E-mail: gauravgovil@yahoo.co.in;

Keywords: Pelvis; symphysis; radiography; ankylosing spondylitis; spondylitis; clinic-radiological.

ABBREVIATION

MRI : Magnetic Resonance Imaging

1. INTRODUCTION

Pubic symphysis frequently presents with painful aetiology due to inflammation. Commonly pelvic instability or pubic inflammation presents with significant discomfort and pain along the pubic region [1]. An asymptomatic incidental presentation of symphyseal affection have been noticed very infrequently or occasionally on imaging of the pelvis for other reasons [2]. Ankylosis of pubic symphysis presents commonly due to either a developmental, arthritic or post-surgical fusion aetiology [3]. An idiopathic aetiology for a fused symphysis pubis is rare and it has been very infrequently detailed.

The incidental radiographic finding in a trauma patient required a clinical re-evaluation with detailed thorough history taking and examination to assess for the possible aetiological factors. We present the image to discuss the significance of history taking for the assessment of a clinico-radiological mismatch. We also highlight the clinical approach to a fused pubic symphyseal affection.

2. CASE REPORT

A 62-year-old male, of South-east Asian origin, well-built, businessman by profession, presented to emergency with history of road traffic accident while walking involving a hit by a two-wheeler.

Injuries were sustained to head, face and pelvic region. After clinical assessment, radiological tests were requested. Computed tomography and plain x-ray were preferred investigations for head and pelvis, respectively. The computer tomographic imaging of head showed haemorrhages in left basal ganglia and corona radiata with surrounding oedema and mild compression of left lateral ventricle. A trauma protocol assessment with a routine pelvis radiography revealed an incidental ankylosis of pubic symphysis. On clinical examination, there were no signs of pelvic or limb trauma. An active straight leg-raising test was possible and no motor sensory deficit was present. Pelvic compression test was negative and there was no focal tenderness at symphysis. After primary stabilization of vitals, a secondary clinical evaluation revealed no past history of pain along the pelvic region. There was negative history related to hip, pelvis and sacroiliac joint suggestive of any inflammatory, arthritic, degenerative or infective pathology as a pre-disposing factor leading to pubic symphysis ankylosis. There was no history of any pelvis related surgical intervention. Symptoms suggestive of pelvic instability could not be ascertained [4].

Initial management required intensive care monitoring for head injury. Radiologically, the fusion of pubic symphysis presented with smooth symphyseal contours with minimal sclerosis but without any evidence of irregularity, osteopenia or any surgical intervention [Fig. 1].



Fig. 1. Anteroposterior radiograph of pelvis with ankylosed pubic symphysis

There was no developmental disparity of pelvic bony architecture. The absence of inter-posed symphyseal fibro-cartilaginous disc was evident. A magnetic resonance imaging (MRI) was suggested for further evaluation of pelvic pathology. However, patient did not consent and further imaging was deferred in view of the asymptomatic nature of the clinico-radiological mismatch presentation.

3. DISCUSSION

Symphysis pubis, a non-synovial fibrocartilaginous joint may present with painful pathologies of varied aetiology [1,2]. A thorough history taking will help to identify the possible aetiology. History related to trauma, fever, multiple joint pain and surgical intervention will guide regarding the probable provisional cause of fusion. The clinical examination should determine the exact site of pain. The duration of symptom indicates an acute or chronic pathology affection. The familial history identifies uncommon cause of developmental disorders.

An ankylosed pubis may result due to retro pubic infection. According to the theory, by Coventry and Mitchell, the retro pubic infection may cause venous stasis, thrombosis and eventual avascular necrosis [4,5]. In females, the common aetiology for infection includes repetitive trauma from multiple pregnancies, pelvic surgeries or pelvic sepsis presenting as osteitis pubis of inflammatory aetiology [1,3]. Osteitis pubis in males commonly affects sportspersons and athletes [1]. Tubercular and bacterial affection may present with severe inflammatory changes of symphysis pubis and infrequently septic arthritis [1,2].

The progression of the radiological changes to induce ankylosis have been hypothesised due to "backfill" by new tissue of bone marrow intensity [2,6]. Symphyseal erosions, enthesitis, widening or degeneration may present with arthritic affections related to rheumatoid, osteoarthritis or ankylosing spondylitis [1,2]. Bridging or fusion of the pubic symphysis can also be associated with various systemic and local causes. This may include fluorosis, ochronosis, post-traumatic, post-radiation therapy and myositis ossificans [7].

The symphyseal affection can be evaluated further with certain biochemical and imaging modalities based on the probable causes.

Estimation of acute phase reactants, C-reactive protein, differential leucocyte count, erythrocyte sedimentation rate, pro-calcitonin may delineate and differentiate the inflammatory aetiology. Serological markers including rheumatoid factor, anti-CCP antibody levels, HLA-B27 estimation may also guide regarding the possible arthritic affection. On radiographs, inflammation commonly presents with symphyseal joint space erosions, sclerosis, widening or obliteration [8]. Computerised tomographic imaging may be helpful to evaluate bony involvement and affections due to trauma and tumour [1]. MRI of the pubic symphyseal region may reveal an active inflammatory change in cases related to spondyloarthropathy affections in the young and elderly age group [3]. MRI evaluation may present with subchondral marrow oedema and peri-symphyseal soft tissue oedema in association with bony changes in the affections related to inflammatory aetiology [1]. The clinicians should interpret the MRI findings in consideration to the age of presentation, intensity and onset of symptoms, and relevant history for any repetitive trauma [9]. MRI presents with reliable findings with high sensitivity to differentiate between the acute or chronic changes involving the pubic symphysis [10]. MRI findings of bone marrow oedema with an associated periarticular oedema in the soft tissue and muscles have poor clinical long-term outcomes [11].

The symphyseal affection normally responds to anti-inflammatory medications. In majority, the treatment with conservative measures may resolve the symptoms of pubic pain. However, when conservative treatment fails, surgical management may be required. The clinically significant outcome improvement can be prognosticated based on the clinical correlation to imaging findings [10]. Surgical option includes wedge resection of inflamed symphysis with bone grafting and fixation with plates to resolve the symptoms [4,8]. Curettage, wide resection or arthrodesis can also be done [8].

4. CONCLUSION

An asymptomatic ankylosed symphysis pubis in a case of trauma may present rarely. The radiological presentation with a clinical mismatch may present challenges for the trauma surgeon and the emergency clinicians. A thorough history taking and awareness of the differential diagnosis will alleviate the diagnostic dilemma and guide in the management.

DATA AVAILABILITY

All data generated or analysed during this study are included in this article. Further enquiries can be directed to the corresponding author.

CONSENT

A written informed consent was taken from the patient for the use of data for publication. The patient understands that name and initials will not be published and due efforts will be made to conceal the identity, but anonymity cannot be guaranteed.

ETHIC APPROVAL

Ethical approval was not required for this study in accordance with local guidelines.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Budak MJ, Oliver TB. There's a hole in my symphysis -- a review of disorders causing widening, erosion, and destruction of the symphysis pubis. *Clin Radiol*. 2013;68(2):173-80. DOI: 10.1016/j.crad.2012.03.021.
2. Koo BS, Song Y, Joo KB, Lee S, Kim TH. Radiologic Changes in the Symphysis Pubis of Male Patients with Ankylosing Spondylitis. *J Rheumatol*. 2016;43(2):330-4. DOI: 10.3899/jrheum.150711.
3. Kang Y, Ahn JM, Lee E, Lee JW, Kang HS. Active inflammatory changes around the pubic symphysis in patients with axial spondyloarthritis: Magnetic resonance imaging characteristics and association with clinical factors. *Eur J Radiol*. 2020;124:108802. DOI: 10.1016/j.ejrad.2019.108802.
4. Harris NH. Lesions of the symphysis pubis in women. *Br Med J*. 1974;4(5938):209-11. DOI: 10.1136/bmj.4.5938.209.
5. Coventry MB, Mitchell WC. Osteitis Pubis. *JAMA*. 1961;178(9):898-905. DOI:10.1001/jama.1961.03040480028006
6. Maksymowych WP, Wichuk S, Chiowchanwisawakit P, Lambert RG, Pedersen SJ. Fat metaplasia and backfill are key intermediaries in the development of sacroiliac joint ankylosis in patients with ankylosing spondylitis. *Arthritis Rheumatol*. 2014;66:2958-67.
7. Mudgal P, Knipe H. Bridging of the pubic symphysis (differential). Reference article, Radiopaedia.org. Accessed on 13 Oct 2021 Available: <https://radiopaedia.org/articles/26990>
8. Nasrallah K, Jammal M, Khoury A, Liebergall M. Adult female patient with osteitis pubis and pelvic instability requiring surgery: A case report. *Trauma Case Rep*. 2020;30:100357. DOI: 10.1016/j.tcr.2020.100357.
9. Thorborg K, Reiman MP, Weir A, Kemp JL, Serner A, Mosler AB, Hölmich P. Clinical Examination, Diagnostic Imaging, and Testing of Athletes With Groin Pain: An Evidence-Based Approach to Effective Management. *J Orthop Sports Phys Ther*. 2018;48(4):239-249. DOI: 10.2519/jospt.2018.7850.
10. Krishnamoorthy VP, Kunze KN, Beck EC, Cancienne JM, O'Keefe LS, Ayeni OR, Nho SJ. Radiographic Prevalence of Symphysis Pubis Abnormalities and Clinical Outcomes in Patients with Femoroacetabular Impingement Syndrome. *The American Journal of Sports Medicine*. 2019;47(6):1467-1472. DOI:10.1177/0363546519837203.
11. Gaudino F, Spira D, Bangert Y, Ott H, Beomonte Zobel B, Kauczor HU, Weber MA. Osteitis pubis in professional football players: MRI findings and correlation with clinical outcome. *Eur J Radiol*. 2017;9(94):46-52. DOI:10.1016/j.ejrad.2017.07.009.

© 2021 Tomar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/81916>