

## Pullback Absorbing Set for the Stochastic Lattice Selkov Equations

Li Hongyan<sup>1\*</sup>

<sup>1</sup>College of Management, Shanghai University of Engineering Science, Shanghai 201620, China.

### Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

### Article Information

DOI: 10.9734/ARJOM/2019/v14i130116

Editor(s):

(1) Dr. Radoslaw Jedynak, Department of Computer Science and Mathematics, Kazimierz Pulaski University of Technology and Humanities, Poland.

Reviewers:

(1) Olumide Adesina, Olabisi Onabanjo University, Nigeria.

(2) Joshua Otonritse Okor, Landmark University, Nigeria.

(3) Abdullah Sonmezoglu, Yozgat Bozok University, Turkey.

(4) Sie Long Kek, Universiti Tun Hussein Onn Malaysia, Malaysia.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48970>

Received: 15 March 2019

Accepted: 21 May 2019

Published: 29 May 2019

Original Research Article

## Abstract

**Aims/ Objectives:** To prove the existence of a pullback Absorbing set.

**Study Design:** Ornstein-Uhlenbeck process.

**Place and Duration of Study:** College of Management, Shanghai University of Engineering Science.

**Methodology:** A transformation of addition involved with an Ornstein-Uhlenbeck process is used.

**Results:** In this paper, pullback absorbing property for the stochastic reversible Selkov system in an infinite lattice with additive noises is proved.

*Keywords:* Pullback absorbing set; additive noise; Selkov system.

**2010 Mathematics Subject Classification:** 34F05

\*Corresponding author: E-mail: hongyanlishu@163.com;

## 1 Introduction

In this paper, we consider the stochastic lattice Selkov system with the cubic nonlinearity and additive white noises on an infinite lattice as follows:


$$\begin{cases} du_i = [d_1(u_{i+1} - 2u_i + u_{i-1}) - a_1u_i + b_1u_i^2v_i - b_2u_i^3 + f_{1i}]dt + \alpha_i dw_i, & i \in \mathbb{Z}, t > 0, \\ dv_i = [d_2(v_{i+1} - 2v_i + v_{i-1}) - a_2v_i - b_1u_i^2v_i + b_2u_i^3 + f_{2i}]dt + \alpha_i dw_i, & i \in \mathbb{Z}, t > 0, \end{cases} \quad (1.1)$$

with initial conditions

$$u_i(0) = u_{i,0}, \quad v_i(0) = v_{i,0}, \quad i \in \mathbb{Z}, \quad (1.2)$$

where  $\mathbb{Z}$  denotes the integer set,  $u = (u_i)_{i \in \mathbb{Z}} \in \ell^2$ ,  $v = (v_i)_{i \in \mathbb{Z}} \in \ell^2$ ,  $d_1, d_2, a_1, a_2, b_1, b_2$  are positive constants,  $\alpha = (\alpha_i)_{i \in \mathbb{Z}} \in \ell^2$ ,  $\{w_i | i \in \mathbb{Z}\}$  is independent Brownian motions.

The reversible Selkov model is derived from a set of the two reversible chemical reactions:



The original Selkov model corresponds to the two irreversible reactions, where the product  $Q$  is an inert product. Let  $u_t$  and  $v_t$  are respectively the concentrations of the reactants  $B$  and  $A$ , Equation (1.1) can be regarded as a Selkov system (see [1]) on  $\mathbb{R}$ :

$$\begin{cases} u_t = d_1 \Delta u - a_1 u + b_1 u^2 v - b_2 u^3 + f_1 + \alpha w_t, \\ v_t = d_2 \Delta v - a_2 v - b_1 u^2 v + b_2 u^3 + f_2 + \alpha w_t. \end{cases} \quad (1.3)$$

Here  $\Delta$  is a Laplacian,  $w_t$  is the white noise to the respective components. We have obtained the random dynamical system, see [2]. Pullback absorbing property is very important to describe the long-time behavior of the equations for the mathematics and physics, especially, to prove the existence of random attractor. Therefore, in this paper, we prove the pullback absorbing property for the Selkov equations (1.1).

### 1.1 Preliminaries

In this section, we introduce the relevant definitions of absorbing property, which are taken from [3], [4], [5], [6].

Assume  $X$  is a separable Banach space. For random parameters, we choose the standard probability space  $(\Omega, \mathcal{F}, P)$  where  $\Omega = \{\omega \in C(\cdot) : \omega(0) = 0\}$ ,  $\mathcal{F}$  is the Borel  $\sigma$ -algebra induced by the compact open topology of  $\Omega$ , and  $P$  is the Wiener measure on  $(\Omega, \mathcal{F})$ .

**Definition 2.1.** If  $\theta : \mathbb{R} \times \Omega \rightarrow \Omega$  is  $(\mathcal{B}(\mathbb{R}) \times \mathcal{F}, \mathcal{F})$  measurable, and

$$\theta_0 = I, \theta_{s+t} = \theta_s \circ \theta_t, \quad \forall s, t \in \mathbb{R},$$

$$\theta_t P = P, \quad \forall t \in \mathbb{R},$$

then  $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$  is called a metric dynamical system.

**Definition 2.2.** If a mapping

$$\psi : \mathbb{R}^+ \times \Omega \times X \rightarrow X, \quad (t, \omega, x) \mapsto \psi(t, \omega, x),$$

is  $(\mathcal{B}^+(\mathbb{R}) \times \mathcal{F} \times \mathcal{B}(X), \mathcal{B}(X))$ -measurable and satisfies, for every  $\omega \in \Omega$ ,

- (i)  $\psi(0, \omega, \cdot)$  is the identity on  $X$ ;

- (ii) Cocycle property:  $\psi(t + s, \omega, \cdot) = \psi(t, \theta_s \omega, \psi(s, \omega, \cdot))$  for all  $t, s \in \mathbb{R}^+$ ;
- (iii)  $\psi(\cdot, \omega, \cdot) : \mathbb{R}^+ \times X \rightarrow X$  is strongly continuous.

Then the map  $\psi$  is a continuous random dynamical system on  $X$  over a metric dynamical system  $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$ .

**Definition 2.3.** If for every  $\omega \in \Omega$  and a random bounded set  $D(\omega) \subset X$ ,

$$\lim_{t \rightarrow \infty} e^{\gamma t} d(D(\theta_{-t} \omega)) = 0 \text{ for all } \gamma > 0,$$

where  $d(D) = \sup_{x \in D} \|x\|_X$ . Then  $D(\omega) \subset X$  is called tempered with respect to  $(\theta_t)_{t \in \mathbb{R}}$ .

**Definition 2.4.** A random set  $J(\omega)$  is called a pullback absorbing set in  $\mathcal{D}$ , if for all  $D \in \mathcal{D}$  and every  $\omega \in \Omega$ , there exists a  $t_D(\omega) > 0$  such that

$$\psi(t, \theta_{-t} \omega, D(\theta_{-t} \omega)) \subset J(\omega), \forall t \geq t_D(\omega).$$

Where  $\mathcal{D}$  is a collection of random sets of  $X$ .

## 2 Ornstein-Uhlenbeck Process

Let  $\iota^2 = \{u = (u_i)_{i \in \mathbb{Z}}, u_i \in \mathbb{R} : \sum_{i \in \mathbb{Z}} |u_i|^2 < +\infty\}$ , with the inner product and norm as follows:

$$\langle u, v \rangle = \sum_{i \in \mathbb{Z}} u_i v_i, \quad \|u\|^2 = \langle u, u \rangle, \quad u = (u_i)_{i \in \mathbb{Z}}, v = (v_i)_{i \in \mathbb{Z}} \in \iota^2.$$

Then  $\iota^2 = (\iota^2, \langle \cdot, \cdot \rangle, \|\cdot\|)$  is a Hilbert space. Set  $E = \iota^2 \times \iota^2$  be the product Hilbert space. In view of the cubic term  $\pm u^2 v, \pm u^3$ , we need  $u \in \iota^6, v \in \iota^6$  to make (1.1) hold in  $\iota^2$ .

To convert the stochastic equation to a deterministic one with random parameters, we introduce an Ornstein-Uhlenbeck process (O-U process) (see [7]) in  $\iota^2$  on  $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$  given by the Wiener process:

$$y(\theta_t \omega) = -(a_1 + a_2) \int_{-\infty}^0 e^{(a_1 + a_2)s} (\theta_t \omega)(s) ds, \quad t \in \mathbb{R}, \quad \omega \in \Omega,$$

and  $y$  solve the following Itô equations respectively:

$$dy + (a_1 + a_2)y dt = dw(t), \quad t > 0.$$

There exists a  $\theta_t$ -invariant set  $\Omega' \subset \Omega$  of full  $P$  measure such that

- (1) the mappings  $s \rightarrow y(\theta_s \omega)$ , is continuous for each  $\omega \in \Omega$ ;
- (2) the random variables  $\|y(\theta_t \omega)\|$  is tempered.

To transform (1.3) into pathwise equations, Denote

$$\tilde{u}(t) = u(t) - y(\theta_t \omega), \quad \tilde{v}(t) = v(t) - y(\theta_t \omega).$$

From (1.3), we have

$$\begin{cases} \tilde{u}_t = -d_1 A(\tilde{u} + y(\theta_t \omega)) - a_1 \tilde{u} + a_2 y(\theta_t \omega) + b_1 (\tilde{u} + y(\theta_t \omega))^2 (\tilde{v} + y(\theta_t \omega)) \\ \quad - b_2 (\tilde{u} + y(\theta_t \omega))^3 + f_1 \\ \tilde{v}_t = -d_2 A(\tilde{v} + y(\theta_t \omega)) - a_2 \tilde{v} + a_1 y(\theta_t \omega) - b_1 (\tilde{u} + y(\theta_t \omega))^2 (\tilde{v} + y(\theta_t \omega)) \\ \quad + b_2 (\tilde{u} + y(\theta_t \omega))^3 + f_2 \end{cases} \quad (2.1)$$

with the initial value condition

$$\tilde{u}(0, \omega, \tilde{u}_0) = \tilde{u}_0(\omega) = u_0 - y(\omega), \quad \tilde{v}(0, \omega, \tilde{v}_0) = \tilde{v}_0(\omega) = v_0 - y(\omega).$$

### 3 Pullback Absorbing Property

**Lemma 4.1.** *There exists a  $\theta_t$ -invariant set  $\Omega' \subset \Omega$  of full  $P$  measure and an absorbing random set  $J(\omega), \omega \in \Omega', \forall D \in \mathcal{D}$  and  $\forall \omega \in \Omega'$ , there exists  $T_D(\omega) > 0$  such that*

$$\psi(t, \theta_{-t}\omega, D(\theta_{-t}\omega)) \subset J(\omega) \quad \forall t \geq T_D(\omega).$$

Moreover,  $J \in \mathcal{D}$ .

**Proof.** Taking the inner product to (2.1) with  $(\tilde{u}, \tilde{v})^T$  in  $E$ , we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\tilde{u}\|^2 &= -d_1 \langle A\tilde{u}, \tilde{u} \rangle - d_1 \langle Ay(\theta_t\omega), \tilde{u} \rangle - a_1 \|\tilde{u}\|^2 + b_1 \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{v} + y(\theta_t\omega)), \tilde{u} \rangle \\ &\quad - b_2 \langle (\tilde{u} + y(\theta_t\omega))^3, \tilde{u} \rangle + \langle f_1, \tilde{u} \rangle + a_2 \langle y(\theta_t\omega), \tilde{u} \rangle, \\ \frac{1}{2} \frac{d}{dt} \|\tilde{v}\|^2 &= -d_2 \langle A\tilde{v}, \tilde{v} \rangle - d_2 \langle Ay(\theta_t\omega), \tilde{v} \rangle - a_2 \|\tilde{v}\|^2 - b_1 \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{v} + y(\theta_t\omega)), \tilde{v} \rangle \\ &\quad + b_2 \langle (\tilde{u} + y(\theta_t\omega))^3, \tilde{v} \rangle + \langle f_2, \tilde{v} \rangle + a_1 \langle y(\theta_t\omega), \tilde{v} \rangle. \end{aligned} \quad (3.1)$$

Summing the two equations up, we have

$$\begin{aligned} &\frac{d}{dt} [\|\tilde{u}\|^2 + \|\tilde{v}\|^2] + 2d_1 \langle A\tilde{u}, \tilde{u} \rangle + 2d_2 \langle A\tilde{v}, \tilde{v} \rangle + 2a_1 \|\tilde{u}\|^2 + 2a_2 \|\tilde{v}\|^2 \\ &= -2d_1 \langle Ay(\theta_t\omega), \tilde{u} \rangle - 2d_2 \langle Ay(\theta_t\omega), \tilde{v} \rangle + 2\langle f_1, \tilde{u} \rangle + 2\langle f_2, \tilde{v} \rangle + 2a_2 \langle y(\theta_t\omega), \tilde{u} \rangle + 2a_1 \langle y(\theta_t\omega), \tilde{v} \rangle \\ &\quad + 2b_1 \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{v} + y(\theta_t\omega)), \tilde{u} \rangle - 2b_1 \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{v} + y(\theta_t\omega)), \tilde{v} \rangle \\ &\quad - 2b_2 \langle (\tilde{u} + y(\theta_t\omega))^3, \tilde{u} \rangle + 2b_2 \langle (\tilde{u} + y(\theta_t\omega))^3, \tilde{v} \rangle. \end{aligned} \quad (3.2)$$

Then we have

$$\begin{aligned} &2b_1 \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{v} + y(\theta_t\omega)), \tilde{u} \rangle - 2b_1 \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{v} + y(\theta_t\omega)), \tilde{v} \rangle \\ &\quad - 2b_2 \langle (\tilde{u} + y(\theta_t\omega))^3, \tilde{u} \rangle + 2b_2 \langle (\tilde{u} + y(\theta_t\omega))^3, \tilde{v} \rangle \\ &= 2b_1 \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{v} + y(\theta_t\omega)), \tilde{u} - \tilde{v} \rangle - 2b_2 \langle (\tilde{u} + y(\theta_t\omega))^3, \tilde{u} - \tilde{v} \rangle \\ &\leq 2 \max\{b_1, b_2\} \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{v} + y(\theta_t\omega)) - \tilde{u} - y(\theta_t\omega), \tilde{u} - \tilde{v} \rangle \\ &= -2 \max\{b_1, b_2\} \langle (\tilde{u} + y(\theta_t\omega))^2 (\tilde{u} - \tilde{v}), \tilde{u} - \tilde{v} \rangle \\ &= -2 \sum_{i \in \mathbb{Z}} (\tilde{u}_i + y_i(\theta_t\omega))^2 (\tilde{u}_i - \tilde{v}_i)^2 \leq 0. \end{aligned} \quad (3.3)$$

By Young's inequality in [8], we have the following estimate

$$-2d_1 \langle Ay(\theta_t\omega), \tilde{u} \rangle \leq \frac{a_1}{3} \|\tilde{u}\|^2 + \frac{3d_1^2}{a_1} \|Ay(\theta_t\omega)\|^2, \quad (3.4)$$

$$-2d_2 \langle Ay(\theta_t\omega), \tilde{v} \rangle \leq \frac{a_2}{3} \|\tilde{v}\|^2 + \frac{3d_2^2}{a_2} \|Ay(\theta_t\omega)\|^2, \quad (3.5)$$

$$2a_2 \langle y(\theta_t\omega), \tilde{u} \rangle \leq \frac{a_1}{3} \|\tilde{u}\|^2 + \frac{3a_2^2}{a_1} \|y(\theta_t\omega)\|^2, \quad (3.6)$$

$$2a_1 \langle y(\theta_t\omega), \tilde{v} \rangle \leq \frac{a_2}{3} \|\tilde{v}\|^2 + \frac{3a_1^2}{a_2} \|y(\theta_t\omega)\|^2, \quad (3.7)$$

$$2\langle f_1, \tilde{u} \rangle \leq \frac{a_1}{3} \|\tilde{u}\|^2 + \frac{3}{a_1} \|f_1\|^2, \quad (3.8)$$

$$2\langle f_2, \tilde{v} \rangle \leq \frac{a_2}{3} \|\tilde{v}\|^2 + \frac{3}{a_2} \|f_2\|^2. \quad (3.9)$$

By (3.2)-(4.9), we obtain that

$$\begin{aligned}
 & \frac{d}{dt} [\|\tilde{u}\|^2 + \|\tilde{v}\|^2] + 2d_1 \langle A\tilde{u}, \tilde{u} \rangle + 2d_2 \langle A\tilde{v}, \tilde{v} \rangle + 2a_1 \|\tilde{u}\|^2 + 2a_2 \|\tilde{v}\|^2 \\
 \leq & \frac{a_1}{3} \|\tilde{u}\|^2 + \frac{3d_1^2}{a_1} \|Ay(\theta_t\omega)\|^2 + \frac{a_2}{3} \|\tilde{v}\|^2 + \frac{3d_2^2}{a_2} \|Ay(\theta_t\omega)\|^2 + \frac{3}{a_1} \|f_1\|^2 + \frac{a_1}{3} \|\tilde{u}\|^2 \\
 & + \frac{3}{a_2} \|f_2\|^2 + \frac{a_2}{3} \|\tilde{v}\|^2 + \frac{3a_2^2}{a_1} \|y(\theta_t\omega)\|^2 + \frac{a_1}{3} \|\tilde{u}\|^2 + \frac{3a_1^2}{a_2} \|y(\theta_t\omega)\|^2 + \frac{a_2}{3} \|\tilde{v}\|^2 \\
 = & a_1 \|\tilde{u}\|^2 + \frac{3d_1^2}{a_1} \|Ay(\theta_t\omega)\|^2 + a_2 \|\tilde{v}\|^2 + \frac{3d_2^2}{a_2} \|Ay(\theta_t\omega)\|^2 + \frac{3}{a_1} \|f_1\|^2 \\
 & + \frac{3}{a_2} \|f_2\|^2 + \frac{3a_2^2}{a_1} \|y(\theta_t\omega)\|^2 + \frac{3a_1^2}{a_2} \|y(\theta_t\omega)\|^2,
 \end{aligned}$$

hence we have,

$$\begin{aligned}
 & \frac{d}{dt} [\|\tilde{u}\|^2 + \|\tilde{v}\|^2] + a_1 \|\tilde{u}\|^2 + a_2 \|\tilde{v}\|^2 \\
 \leq & \frac{3d_1^2}{a_1} \|Ay(\theta_t\omega)\|^2 + \frac{3d_2^2}{a_2} \|Ay(\theta_t\omega)\|^2 + \frac{3}{a_1} \|f_1\|^2 + \frac{3}{a_2} \|f_2\|^2 + \frac{3a_2^2}{a_1} \|y(\theta_t\omega)\|^2 + \frac{3a_1^2}{a_2} \|y(\theta_t\omega)\|^2 \\
 \leq & C_1 \|Ay(\theta_t\omega)\|^2 + C_2 \|y(\theta_t\omega)\|^2 + C_3 (\|f_1\|^2 + \|f_2\|^2) \\
 \leq & C_4 (\|y(\theta_t\omega)\|^2 + \|Ay(\theta_t\omega)\|^2 + \|f_1\|^2 + \|f_2\|^2 + \|f_3\|^2), \tag{3.10}
 \end{aligned}$$

where

$$\begin{aligned}
 C_1 &= \max\left\{\frac{3d_1^2}{a_1}, \frac{3d_2^2}{a_2}\right\}, & C_2 &= \max\left\{\frac{3a_2^2}{a_1}, \frac{3a_1^2}{a_2}\right\}, \\
 C_3 &= \max\left\{\frac{3}{a_1}, \frac{3}{a_2}\right\}, & C_4 &= \max\{C_1, C_2, C_3\}.
 \end{aligned}$$

By Gronwall's inequality in [8], it follows that

$$\begin{aligned}
 & \|\tilde{u}(t, \omega, \tilde{u}_0(\omega))\|^2 + \|\tilde{v}(t, \omega, \tilde{v}_0(\omega))\|^2 \\
 \leq & e^{-\min\{a_1, a_2\}t} [\|\tilde{u}_0(\omega)\|^2 + \|\tilde{v}_0(\omega)\|^2] + \frac{C_4}{\min\{a_1, a_2\}} (\|f_1\|^2 + \|f_2\|^2) \\
 & + C_4 \int_0^t e^{-\min\{a_1, a_2\}(t-s)} (\|y(\theta_s\omega)\|^2 + \|Ay(\theta_s\omega)\|^2) ds. \tag{3.11}
 \end{aligned}$$

Let  $c_1 = \min\{a_1, a_2\}$ . Now that the random variable  $y(\theta_t\omega)$  is tempered and continuous in  $t$ . It follows from Proposition 4.3.3 in [9], there is a tempered function  $j(\omega) > 0$  that satisfies

$$\|y(\theta_t\omega)\|^2 + \|Ay(\theta_t\omega)\|^2 \leq j(\theta_t\omega) \leq j(\omega)e^{\frac{c_1}{2}|t|}. \tag{3.12}$$

Replacing  $\omega$  by  $\theta_{-t}\omega$  in (3.11), by (3.12), we get

$$\begin{aligned}
 & \|\tilde{u}(t, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 + \|\tilde{v}(t, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))\|^2 \\
 \leq & e^{-c_1 t} [\|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t}\omega)\|^2] + \frac{C_4}{c_1} (\|f_1\|^2 + \|f_2\|^2) \\
 & + C_4 \int_0^t e^{-c_1(t-s)} (\|y(\theta_{s-t}\omega)\|^2 + \|Ay(\theta_{s-t}\omega)\|^2) ds \\
 \leq & e^{-c_1 t} [\|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t}\omega)\|^2] + \frac{C_4}{c_1} (\|f_1\|^2 + \|f_2\|^2) \\
 & + C_4 \int_{-t}^0 e^{c_1\tau} (\|y(\theta_\tau\omega)\|^2 + \|Ay(\theta_\tau\omega)\|^2) d\tau \\
 \leq & e^{-c_1 t} [\|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t}\omega)\|^2] + \frac{C_4}{c_1} (\|f_1\|^2 + \|f_2\|^2) + \frac{2C_4 l(\omega)}{c_1}. \tag{3.13}
 \end{aligned}$$

Define  $R^2(\omega) = 2[C_4(\|f_1\|^2 + \|f_2\|^2) + 2C_4l(\omega)]/c_1$ ;  $J(\omega)$  is a tempered function, so  $R(\omega)$  is also tempered.

Define

$$\tilde{J}(\omega) = \{(\tilde{u}, \tilde{v}) \in \ell^2 \times \ell^2, \|\tilde{u}\|^2 + \|\tilde{v}\|^2 \leq R^2(\omega)\}.$$

From Theorem 4.2 in [2],  $\tilde{J}(\omega)$  is an absorbing set for the random dynamical system  $(\tilde{u}(t, \omega, \tilde{u}_0), \tilde{v}(t, \omega, \tilde{v}_0))$ , i.e.,  $\forall D \in \mathcal{D}$  and  $\forall \omega \in \Omega'$ , there exists  $T_D(\omega)$  such that

$$\Phi(t, \theta_{-t}\omega, D(\theta_{-t}\omega)) \subset \tilde{J}(\omega) \quad \text{for } t \geq T_D(\omega).$$

Let

$$J(\omega) = \{(u, v) \in \ell^2 \times \ell^2, \|u\|^2 + \|v\|^2 \leq R_1^2(\omega)\},$$

where

$$R_1^2(\omega) = 2R^2(\omega) + 4\|y(\theta_t\omega)\|^2.$$

since

$$\begin{aligned} & \psi(t, \omega, (u_0, v_0, z_0)) \\ &= \Phi(t, \omega, (u_0 - y(\omega), v_0 - y(\omega))) + (y(\theta_t\omega), y(\theta_t\omega)) \\ &= (\tilde{u}(t, \omega, u_0 - y(\omega)) + y(\theta_t\omega), \tilde{v}(t, \omega, v_0 - y(\omega)) + y(\theta_t\omega)), \end{aligned}$$

so  $J(\omega)$  is an absorbing random set for  $\psi(t, \omega)$  and  $J \in \mathcal{D}$ . The proof of Lemma 4.1 is completed.  $\square$

## Acknowledgement

The paper is supported by Shanghai Natural Science Foundation (16ZR1414000).

## Competing Interests

The author has declared that no competing interests exist.

## References

- [1] Selkov E. Self-oscillations in glycolysis. I. A simple kinetic model. Eur. J. Biochem. 1968;4:79-86.
- [2] Li H. A random dynamical system of the stochastic lattice reversible Selkov equations. Differential Equations and Applications; 2019. submitted
- [3] Bates P, Lisei H, Lu K. Attractors for stochastic lattice dynamical systems. Stochastics and Dynamics. 2006;6:1-21.
- [4] Crauel H, Flandoli F. Attractor for random dynamical systems. Probability Theory and Related Fields. 1994;100:365-393.
- [5] Huang J. The random attractor of stochastic fitz hugh-nagumo equations in an infinite lattice with white noises. Physica D. 2007;233:83-94.
- [6] Li H, Tu J. Random attractors for stochastic lattice reversible Gray-Scott systems with additive noise. Electron. J. Diff. Equ. 2015;2015:1-25.

- [7] Li H. Random attractor of the stochastic lattice reversible selkov equations with additive noises. IEEE 13th International Conference on e-Business Engineering (ICEBE); 2016.
- [8] Témam R. Infinite-dimensional dynamical systems in mechanics and physics. Appl. Math. Sci. Springer-Verlag; 1988.
- [9] Arnold L. Random dynamical systems. Springer Monographs in Mathematics, Springer-Verlag; 1998.

---

©2019 Hongyan; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

**Peer-review history:**

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<http://www.sdiarticle3.com/review-history/48970>